

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 236 719 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(43) Date of publication:
04.09.2002 Bulletin 2002/36

(21) Application number: **00979026.2**

(22) Date of filing: **29.11.2000**

(51) Int Cl.7: **C07D 217/26, C07D 401/12,
C07D 405/12, C07D 413/12,
C07D 413/14, A61K 31/472,
A61K 31/4725, A61P 3/04,
A61P 3/06, A61P 3/10,
A61P 9/10, A61P 29/00**

(86) International application number:
PCT/JP00/08464

(87) International publication number:
WO 01/040192 (07.06.2001 Gazette 2001/23)

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **03.12.1999 JP 34554399
27.09.2000 JP 2000295108**

(71) Applicant: **Kyoto Pharmaceutical Industries, Ltd.
Kyoto-shi, Kyoto 604-8444 (JP)**


(72) Inventors:
• **Matsui, Hiroshi**
Nara-shi, Nara 631-0074 (JP)
• **Kobayashi, Hideo**
Kyoto-shi, Kyoto 616-8075 (JP)

• **Azukizawa, Satoru**
Minami-ku Kyoto-shi, Kyoto 601-8213 (JP)
• **Kaai, Masayasu**
Yamatokoriyama-shi, Nara 639-1134 (JP)
• **Yoshimi, Akihisa**
Takatsuki-shi, Osaka 569-0056 (JP)
• **Shirahase, Hiroaki**
Nagaokakyo-shi, Kyoto 617-0814 (JP)

(74) Representative:
von Kreisler, Alek, Dipl.-Chem. et al
Patentanwälte
von Kreisler-Selting-Werner
Postfach 10 22 41
50462 Köln (DE)

(54) **NOVEL HETEROCYCLIC COMPOUNDS AND SALTS THEREOF AND MEDICINAL USE OF THE SAME**

(57) A heterocyclic compound of the formula [I]



wherein R¹ is hydrogen atom or lower alkyl, R² is hydrogen atom, alkyl optionally having a substituent and the like, R³ is hydrogen atom, lower alkyl and the like, A is a single bond or >N-R⁵ wherein R⁵ is hydrogen atom or lower alkyl, B is lower alkylene, and Y is aryl optionally having a substituent and the like, and a pharmaceutically acceptable salt thereof show a hypoglycemic action, a blood hypolipidemic action, an insulin resistance-improving action and a PPAR activating action, and are useful as a hypoglycemic agent, a hypolipidemic agent, an insulin resistance improver, a therapeutic agent of diabetes, a therapeutic agent of diabetic complication, a glucose tolerance improver, an anti-atherosclerosis agent, an anti-obesity agent, an antiinflammatory agent, an agent for the prophylaxis or treatment of

PPAR-mediated disease and an agent for the prophylaxis or treatment of syndrome X.

Description

TECHNICAL FIELD OF THE INVENTION

[0001] The present invention relates to a novel heterocyclic compound and a pharmaceutically acceptable salt thereof, which have a hypoglycemic action, a blood hypolipidemic action, an insulin resistance-improving action or a PPAR (peroxisome proliferator-activated receptor)-activating action. The present invention also relates to a pharmaceutical composition comprising the above-mentioned novel heterocyclic compound or a pharmaceutically acceptable salt thereof. Furthermore, the present invention relates to a hypoglycemic agent, a hypolipidemic agent, an insulin resistance improver, a therapeutic agent of diabetes, an anti-diabetic complication agent (i.e., a therapeutic agent of diabetic complication), a glucose tolerance improver, an anti-atherosclerosis agent, an anti-obesity agent, an antiinflammatory agent, an agent for the prophylaxis or treatment of PPAR-mediated disease and an agent for the prophylaxis or treatment of syndrome X, all of which comprising the above-mentioned novel heterocyclic compound or a pharmaceutically acceptable salt thereof.

BACKGROUND OF THE INVENTION


[0002] As a therapeutic agent of diabetes, biguanide compounds having, as a main action, an inhibitory action on glucose absorption via the intestinal tract and on glucose release from the liver, sulfonylurea compounds having an accelerating action on insulin secretion as a main action, insulin and the like have been employed. However, biguanide compounds cause lactic acidosis, and sulfonylurea compounds sometimes cause serious hypoglycemia due to their strong hypoglycemic action. Therefore, a due care should be given when in use of these compounds. In recent years, there have been active researches and developments of a therapeutic agent of diabetes, which is free of these defects, with the consequence that various compounds having an insulin resistance-improving action have been found.

[0003] The insulin resistance is important as a cause of non-insulin dependent diabetes mellitus (NIDDM), along with decrease in the insulin secretion. Therefore, the development of a pharmaceutical agent that improves insulin resistance has been desired. Various thiazolidine compounds are known as the agent capable of improving the insulin resistance. As these compounds, for example, 5-[4-[(6-hydroxy-2,5,7,8-tetramethylchroman-2-yl)methoxy]benzyl]-2,4-thiazolidinedione (general name: troglitazone) is described in JP-B-2-31079, 5-[[4-[2-(5-ethyl-pyridin-2-yl)ethoxy]phenyl]methyl]-2,4-thiazolidinedione (general name: pioglitazone) is described in JP-B-5-66956, and 5-[[4-[2-[N-methyl-N-(pyridin-2-yl)amino]ethoxy]phenyl]methyl]-2,4-thiazolidinedione (general name: rosiglitazone) is described in JP-A-1-131169.

[0004] It is therefore an object of the present invention to provide a compound having a hypoglycemic action, a blood hypolipidemic action, an insulin resistance-improving action and a PPAR activating action, which has a structure completely different from that of conventional compounds, thereby to increase the diversity in and to broaden the range of selection from hypoglycemic agents, hypolipidemic agents, insulin resistance improvers, therapeutic agents of diabetes, therapeutic agents of diabetic complications, glucose tolerance improvers, anti-atherosclerosis agents, anti-obesity agents, antiinflammatory agents, agents for the prophylaxis or treatment of PPAR-mediated disease and agents for the prophylaxis or treatment of syndrome X.

DISCLOSURE OF THE INVENTION

[0005] The present inventors have conducted intensive studies in an attempt to solve the above-mentioned problems and found that a heterocyclic compound having a novel structure of the formula [I]



wherein

R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent, alkenyl optionally having a substituent, alkynyl optionally having

a substituent, heterocyclic alkyl optionally having a substituent or -COR⁴

wherein R⁴ is hydrogen atom, alkyl optionally having a substituent, aryl optionally having a substituent, alkenyl optionally having a substituent, arylalkyl optionally having a substituent or heterocyclic residue optionally having a substituent,

R³ is hydrogen atom, lower alkyl or lower alkoxy,

A is a single bond or >N-R⁵

wherein R⁵ is hydrogen atom or lower alkyl,


B is lower alkylene, and

Y is aryl optionally having a substituent or an aromatic heterocyclic residue optionally having a substituent

and a pharmaceutically acceptable salt thereof have a hypoglycemic action, a blood hypolipidemic action, an insulin resistance-improving action and a PPAR activating action, which resulted in the completion of the invention.

[0006] Accordingly, the present invention relates to the following.

[1] A heterocyclic compound of the formula [I]



wherein

R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent, alkenyl optionally having a substituent, alkynyl optionally having a substituent, heterocyclic alkyl optionally having a substituent or -COR⁴ wherein R⁴ is hydrogen atom, alkyl optionally having a substituent, aryl optionally having a substituent, alkenyl optionally having a substituent, arylalkyl optionally having a substituent or heterocyclic residue optionally having a substituent

R³ is hydrogen atom, lower alkyl or lower alkoxy,

A is a single bond or >N-R⁵

wherein R⁵ is hydrogen atom or lower alkyl,

B is lower alkylene, and

Y is aryl optionally having a substituent or an aromatic heterocyclic residue optionally having a substituent (hereinafter to be referred to as

heterocyclic compound [I]), and a pharmaceutically acceptable salt thereof;

[2] the heterocyclic compound of the above-mentioned [1]

wherein, in the formula [I],

R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent or -COR⁴ wherein R⁴ is hydrogen atom, alkyl optionally having a substituent, aryl optionally having a substituent or arylalkyl optionally having a substituent,

R³ is hydrogen atom, lower alkyl or lower alkoxy,

A is a single bond or >N-R⁵ wherein R⁵ is hydrogen atom or lower alkyl,

B is lower alkylene, and

Y is aryl optionally having a substituent or an aromatic heterocyclic residue optionally having a substituent,

and a pharmaceutically acceptable salt thereof;

[3] the heterocyclic compound of the above-mentioned [1]

wherein, in the formula [I],

R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl, cycloalkylalkyl, arylalkyl optionally having a substituent, alkenyl, alkynyl, heterocyclic alkyl or

-COR⁴ wherein R⁴ is alkyl, alkenyl or aryl,

R³ is hydrogen atom or lower alkoxy,

A is a single bond or >N-R⁵ wherein R⁵ is lower alkyl,

B is lower alkylene, and

Y is aryl or an aromatic heterocyclic residue optionally having a substituent,

and a pharmaceutically acceptable salt thereof;

[4] the heterocyclic compound of the above-mentioned [1]

wherein, in the formula [I],

R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl, cycloalkylalkyl, arylalkyl optionally having a substituent or -COR⁴ wherein R⁴ is alkyl or aryl,

R³ is hydrogen atom,

A is a single bond or >N-R⁵ wherein R⁵ is lower alkyl,

B is lower alkylene, and










Y is an aromatic heterocyclic residue optionally having a substituent,

and a pharmaceutically acceptable salt thereof;


[5] the heterocyclic compound of the above-mentioned [1]

wherein, in the formula [I],

Y-A- is



or



wherein

R^A is isopropyl or tert-butyl,

R^B is isopropyl or tert-butyl,

R^C is isopropyl, tert-butyl, phenyl, thiophen-2-yl, 2-methylpropenyl, 3-butenyl, cyclopropyl, 1-butenyl or









2,2-dimethylpropyl,

and a pharmaceutically acceptable salt thereof;


[6] the heterocyclic compound of the above-mentioned [1]

wherein, in the formula [I],

Y-A- is



or



wherein

R^A is isopropyl or tert-butyl,

R^B is isopropyl or tert-butyl,




R^C is isopropyl, tert-butyl, phenyl, thiophen-2-yl, 2-methylpropenyl or 3-butenyl,


and a pharmaceutically acceptable salt thereof;

[7] the heterocyclic compound of the above-mentioned [1]

wherein, in the formula [I],


Y-A- is





and a pharmaceutically acceptable salt thereof;
 [8] the heterocyclic compound of the above-mentioned [1]
 wherein, in the formula [I],

Y-A- is



and a pharmaceutically acceptable salt thereof;
 [9] the heterocyclic compound of the above-mentioned [1]
 wherein the heterocyclic compound of the formula [I] is any of the following compounds (1) to (67):

- (1) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (2) 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (3) 2-acetyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (4) 2-methyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (5) 2-hexanoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (6) 2-hexyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (7) 2-isobutyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (8) 2-cyclohexylmethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (9) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(3-phenylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (10) 2-benzoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (11) 2-benzyl-7-[2-(N-methyl-N-(pyridin-2-yl)amino)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (12) 2-benzyl-7-[2-(5-ethyl-pyridin-2-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (13) 2-benzyl-7-[2-(indolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (14) ethyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (15) methyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (16) 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (17) ethyl 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (18) 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (19) ethyl 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,

- (20) 2-benzyl-7-[2-(6-carboxyindolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (21) 2-(4-fluorobenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (22) 2-(2,2-dimethylpropionyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (23) 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (24) 2-benzyl-7-[2-(5-methyl-2-tert-butyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (25) 2-benzyl-7-[2-(5-methyl-2-(thiophen-2-yl)oxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (26) 2-benzyl-7-[2-(5-methyl-2-isopropylloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (27) 2-butyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (28) 2-benzyl-7-[2-(5-methyl-2-(2-methylpropenyl)oxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (29) 2-benzyl-7-[2-(2-(3-butenyl)-5-methyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (30) 2-allyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (31) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(2-propynyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (32) 2-(2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (33) 2-benzyl-7-[(indolin-3-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (34) 2-(3-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (35) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-pentanoyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (36) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(4-pentenoyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (37) 2-(3-methyl-2-butenoyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (38) 2-(3,3-dimethylbutyryl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (39) 2-benzyl-7-methoxy-6-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3RS)-carboxylic acid,
- (40) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(pyridin-2-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (41) 2-benzyl-7-(3-methyl-3-phenylbutoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (42) 2-benzyl-7-(3,3-dimethyl-4-phenylbutoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (43) 2-benzyl-7-(2-isopropylbenzoxazol-6-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (44) 2-benzyl-7-(2-tert-butylbenzoxazol-6-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (45) 2-benzyl-7-(2-tert-butylbenzoxazol-5-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (46) 7-(2-tert-butylbenzoxazol-6-yl)methoxy-2-(2,2-dimethylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (47) 2-benzyl-7-(2-isopropylbenzoxazol-5-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (48) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(pyridin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (49) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-[(pyridin-2-yl)carbonyl]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (50) methyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (51) 2-benzyl-7-[2-(2-cyclopropyl-5-methyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (52) 2-(3-methyl-2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (53) 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-tert-butyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(54) 2-benzyl-7-[2-(1-butenyl)-5-methyloxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(55) 2-benzyl-7-[2-(2,2-dimethylpropyl)-5-methyloxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(56) ethyl 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,

(57) 7-(benzofran-2-ylmethoxy)-2-benzyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(58) 2-isobutyryl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(59) 7-[2-(benzofran-2-yl)ethoxy]-2-benzyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(60) 7-[2-(5-ethylpyridin-2-yl)ethoxy]-2-hexanoyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(61) 2-carboxymethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(62) 2-[3-(methoxycarbonyl)propionyl]-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(63) 2-[3-(ethoxycarbonyl)propyl]-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(64) 2-benzyl-6-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3RS)-carboxylic acid,

(65) 2-(3-acetylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(66) 2-(2-acetylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid, and

(67) 2-benzyl-7-[(5-methyl-2-phenyloxazol-4-yl)methoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid, and a pharmaceutically acceptable salt thereof;

[10] the heterocyclic compound of the above-mentioned [9]

wherein the heterocyclic compound of the formula [I] is any of the above-mentioned compounds (1) to (47), and a pharmaceutically acceptable salt thereof;

[11] the heterocyclic compound of the above-mentioned [9]

wherein the heterocyclic compound of the formula [I] is any of the above-mentioned compounds (1) to (21), and a pharmaceutically acceptable salt thereof;

[12] a pharmaceutical composition comprising the heterocyclic compound of any of the above-mentioned [1] to [11] or a pharmaceutically acceptable salt thereof;

[13] a pharmaceutical agent comprising the heterocyclic compound of any of the above-mentioned [1] to [11] or a pharmaceutically acceptable salt thereof, which is selected from the group consisting of a hypoglycemic agent, a hypolipidemic agent, an insulin resistance improver, a therapeutic agent of diabetes, a therapeutic agent of diabetic complication, a glucose tolerance improver, an anti-atherosclerosis agent, an anti-obesity agent, an antiinflammatory agent, an agent for the prophylaxis or treatment of PPAR-mediated disease and an agent for the prophylaxis or treatment of syndrome X;

[14] a hypoglycemic agent comprising the heterocyclic compound of any of the above-mentioned [1] to [11] or a pharmaceutically acceptable salt thereof;


[15] a hypolipidemic agent comprising the heterocyclic compound of any of the above-mentioned [1] to [11] or a pharmaceutically acceptable salt thereof;

[16] an insulin resistance improver comprising the heterocyclic compound of any of the above-mentioned [1] to [11] or a pharmaceutically acceptable salt thereof;

[17] a therapeutic agent of diabetic complications, comprising the heterocyclic compound of any of the above-mentioned [1] to [11] or a pharmaceutically acceptable salt thereof; and

[18] a therapeutic agent of diabetes, comprising the heterocyclic compound of any of the above-mentioned [1] to [11] or a pharmaceutically acceptable salt thereof.

[0007] The novel heterocyclic compound of the formula [I]



wherein

R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent, alkenyl optionally having a substituent, alkynyl optionally having a substituent, heterocyclic alkyl optionally having a substituent or -COR⁴

wherein R⁴ is hydrogen atom, alkyl optionally having a substituent, aryl optionally having a substituent, alkenyl optionally having a substituent, arylalkyl optionally having a substituent or heterocyclic residue optionally having a substituent,

R³ is hydrogen atom, lower alkyl or lower alkoxy,

A is a single bond or >N-R⁵
wherein R⁵ is hydrogen atom or lower alkyl,

B is lower alkylene, and

Y is aryl optionally having a substituent or an aromatic heterocyclic residue optionally having a substituent

and a pharmaceutically acceptable salt thereof have a hypoglycemic action, a blood hypolipidemic action, an insulin resistance-improving action and a PPAR-activating action.

[0008] The alkoxycarbonyl in the present invention preferably has 2 to 5 carbon atoms. Examples thereof include methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl and the like.

[0009] The lower alkyl at R¹, R³ and R⁵ is preferably straight chain or branched chain alkyl having 1 to 6 carbon atoms, which is exemplified by methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, hexyl and the like, preferably methyl, ethyl, propyl and isopropyl.

[0010] The lower alkoxy at R³ is preferably straight chain or branched chain alkoxy having 1 to 6 carbon atoms, which is exemplified by methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy, isopentyloxy, neopentyloxy, hexyloxy and the like, preferably methoxy, ethoxy, propoxy and isopropoxy.

[0011] The "alkenyl" in the alkenyl optionally having a substituent at R² and R⁴ is preferably straight chain or branched chain alkenyl having 2 to 6 carbon atoms, which is exemplified by vinyl, 1-propenyl, 2-propenyl, isopropenyl, allyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 3-methyl-2-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl and the like, preferably allyl, 2-butenyl, 3-butenyl, 4-pentenyl, 2-propenyl and 2-methyl-1-propenyl. As the substituent, exemplified are lower alkoxy (as defined for lower alkoxy at R³), hydroxy, carboxy, alkoxycarbonyl, halogen atom (chlorine atom, bromine atom, iodine atom and fluorine atom), nitro, amino and the like. When alkenyl at R² or R⁴ is substituted, the number of the substituent is preferably 1 or 2.

[0012] The "alkynyl" in the alkynyl optionally having a substituent at R² is preferably straight chain or branched chain alkynyl having 2 to 4 carbon atoms, which is exemplified by ethynyl, 1-propynyl, 2-propynyl, 1-methyl-2-propynyl and the like, preferably ethynyl and 2-propynyl. As the substituent, exemplified are lower alkoxy (as defined for lower alkoxy at R³), hydroxy, carboxy, alkoxycarbonyl, halogen atom (chlorine atom, bromine atom, iodine atom and fluorine atom), nitro, amino and the like. When alkynyl at R² is substituted, the number of the substituent is preferably 1 or 2.

[0013] The "alkyl" in the alkyl optionally having a substituent at R² and R⁴ is preferably straight chain or branched chain alkyl having 1 to 8 carbon atoms, which is exemplified by methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl and the like, preferably methyl, ethyl, isobutyl, propyl, hexyl, pentyl and isopropyl. As the substituent, exemplified are lower alkoxy (as defined for lower alkoxy at R³), hydroxy, carboxy, alkoxycarbonyl, halogen atom (chlorine atom, bromine atom, iodine atom and fluorine atom), nitro, amino and the like. When alkyl at R² is substituted, the number of the substituent is preferably 1 or 2.

[0014] The cycloalkyl at R² is preferably cycloalkyl having 3 to 8 carbon atoms, which is exemplified by cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and the like, preferably cyclopropyl, cyclopentyl and cyclohexyl.

[0015] The cycloalkylalkyl at R² is that wherein cycloalkyl moiety is preferably cycloalkyl having 3 to 8 carbon atoms and alkyl moiety is preferably straight chain or branched chain alkyl having 1 to 3 carbon atoms. Examples thereof are

cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cycloheptylmethyl, cyclooctylmethyl, 2-cyclopropylethyl, 2-cyclobutylethyl, 2-cyclopentylethyl, 2-cyclohexylethyl, 2-cycloheptylethyl, 2-cyclooctylethyl, 3-cyclopropylpropyl, 3-cyclobutylpropyl, 3-cyclopentylpropyl, 3-cyclohexylpropyl, 3-cycloheptylpropyl, 3-cyclooctylpropyl, 1-cyclohexylethyl, 1-cyclohexylpropyl, 2-cyclohexylpropyl and the like, preferably cyclohexylmethyl, 2-cyclohexylethyl, cyclopentylmethyl and 2-cyclopentylethyl.

[0016] The aryl in the aryl optionally having a substituent at R², R⁴ and Y is exemplified by phenyl, naphthyl and the like. As the substituent, exemplified are lower alkyl (as defined for lower alkyl at R¹, R³ and R⁵), lower alkoxy (as defined for lower alkoxy at R³), hydroxy, carboxy, alkoxycarbonyl, halogen atom (chlorine atom, bromine atom, iodine atom and fluorine atom), nitro, amino, acyl (e.g., formyl, acetyl, propanoyl etc.) and the like. When aryl at Y is substituted, the number of the substituent is preferably 1 or 2.

[0017] The arylalkyl optionally having a substituent at R² and R⁴ is exemplified by that wherein aryl moiety is preferably phenyl, naphthyl and the like, and alkyl moiety is preferably straight chain or branched chain alkyl having 1 to 3 carbon atoms. Examples of arylalkyl are benzyl, 1-naphthylmethyl, 2-naphthylmethyl, 2-phenylethyl, 2-(1-naphthyl)ethyl, 2-(2-naphthyl)ethyl, 3-phenylpropyl, 3-(1-naphthyl)propyl, 3-(2-naphthyl)propyl, 1-phenylethyl, 2-phenylpropyl, 1-(1-naphthyl)ethyl, 1-(2-naphthyl)ethyl, 1-(1-naphthyl)propyl, 1-(2-naphthyl)propyl, 2-(1-naphthyl)propyl, 2-(2-naphthyl)propyl and the like, preferably benzyl, 3-phenylpropyl, 1-naphthylmethyl and 2-naphthylmethyl. As the substituent, exemplified are lower alkyl (as defined for lower alkyl at R¹, R³ and R⁵), lower alkoxy (as defined for lower alkoxy at R³), hydroxy, carboxy, alkoxycarbonyl, halogen atom (chlorine atom, bromine atom, iodine atom and fluorine atom), nitro, amino, acyl (e.g., formyl, acetyl, propanoyl etc.) and the like, preferably lower alkyl, lower alkoxy, halogen atom and acyl. When arylalkyl at Y is substituted, the number of the substituent is preferably 1 or 2.

[0018] The aromatic heterocycle in the aromatic heterocyclic residue optionally having a substituent at Y is preferably a monocyclic heterocycle or a condensed heterocycle containing at least one hetero atom selected from the group consisting of oxygen atom, nitrogen atom and sulfur atom. The condensed heterocycle in the present invention has two rings and encompasses that having hetero atom(s) on the both rings. Preferable monocyclic heterocycle includes a 5- or 6-membered ring. The heterocycle constituting the condensed heterocycle is preferably a 5- or 6-membered ring. The ring without a hetero atom, which constitutes the condensed heterocycle, is preferably a 5- or 6-membered ring. Examples of the aromatic heterocyclic residue are monocyclic heterocyclic residue such as furyl, thienyl, pyridyl, imidazolyl, pyrazolyl, oxazolyl, isooxazolyl, thiazolyl, triazolyl, thiadiazolyl, oxadiazolyl, pyridazinyl, pyrimidinyl or pyrazinyl and the like; a condensed heterocyclic residue such as indolyl, isoindolyl, indolinyl, isoindolinyl, indazolyl, benzofuranyl, benzothiophenyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, quinolyl, isoquinolyl, benzoxazinyl, benzothiazinyl, furo[2,3-b]pyridyl, thieno[2,3-b]pyridyl, naphthyridinyl, imidazopyridyl, oxazolopyridyl, thiazolopyridyl and the like, with preference given to pyridyl, oxazolyl, indolinyl, benzoxazolyl, thiazolyl, benzothiazolyl, indolyl, quinolyl and benzofuranyl. As the substituent, exemplified are lower alkyl (as defined for lower alkyl at R¹, R³ and R⁵), lower alkoxy (as defined for lower alkoxy at R³), hydroxy, carboxy, alkoxycarbonyl, halogen atom (chlorine atom, bromine atom, iodine atom and fluorine atom), nitro, amino aryl (e.g., phenyl, naphthyl and the like), heterocyclic residue (e.g., thienyl, pyridyl, furyl and the like), alkenyl (as defined for alkenyl at R² and R⁴), cycloalkyl (e.g., cyclopropyl etc.) and the like, preferably aryl, lower alkyl, carboxy, heterocyclic residue, alkenyl and cycloalkyl. When aromatic heterocyclic residue at Y is substituted, the number of the substituent is preferably 1 or 2.

[0019] The lower alkylene at B is preferably straight chain or branched chain alkylene having 1 to 6 carbon atoms, which is exemplified by methylene, ethylene, trimethylene, tetramethylene, pentamethylene, hexamethylene, methylmethylene, 2,2-dimethyltrimethylene, 2-ethyltrimethylene, 1-methyltetramethylene, 2-methyltetramethylene, 3-methyltetramethylene, 3,3-dimethyltrimethylene, 3,3-dimethyltetramethylene and the like preferably ethylene, trimethylene and tetramethylene.

[0020] The heterocyclic moiety in the heterocyclic alkyl optionally having a substituent at R² is as defined for the "aromatic heterocyclic residue" in the "aromatic heterocyclic residue optionally having a substituent" at Y. As the alkyl moiety, exemplified is straight chain or branched chain alkyl having 1 to 3 carbon atoms. The specific examples of the heterocyclic alkyl are 1-pyridylmethyl, 2-pyridylmethyl, 3-pyridylmethyl, 4-pyridylmethyl; 1-(1-pyridyl)ethyl, 1-(2-pyridyl)ethyl, 1-(3-pyridyl)ethyl, 1-(4-pyridyl)ethyl, 2-(1-pyridyl)ethyl, 2-(2-pyridyl)ethyl, 2-(3-pyridyl)ethyl, 2-(4-pyridyl)ethyl; 1-(1-pyridyl)propyl, 1-(2-pyridyl)propyl, 1-(3-pyridyl)propyl, 1-(4-pyridyl)propyl, 2-(1-pyridyl)propyl, 2-(2-pyridyl)propyl, 2-(3-pyridyl)propyl, 2-(4-pyridyl)propyl, 3-(1-pyridyl)propyl, 3-(2-pyridyl)propyl, 3-(3-pyridyl)propyl, 3-(4-pyridyl)propyl; 2-thienylmethyl, 3-thienylmethyl; 1-(2-thienyl)ethyl, 1-(3-thienyl)ethyl, 2-(2-thienyl)ethyl, 2-(3-thienyl)ethyl; 1-(2-thienyl)propyl, 1-(3-thienyl)propyl, 2-(2-thienyl)propyl, 2-(3-thienyl)propyl, 3-(2-thienyl)propyl, 3-(3-thienyl)propyl; and the like. The heterocyclic alkyl may be substituted at the heterocyclic moiety. As the substituent, exemplified are lower alkyl (as defined for lower alkyl at R¹, R³ and R⁵), lower alkoxy (as defined for lower alkoxy at R³), hydroxy, carboxy, alkoxycarbonyl, halogen atom (chlorine atom, bromine atom, iodine atom and fluorine atom), nitro, amino and the like. When the heterocyclic moiety is substituted, the number of the substituent is preferably 1 or 2.

[0021] With regard to the heterocyclic residue optionally having a substituent at R⁴, the heterocyclic moiety is the same as the "aromatic heterocyclic residue" in the "aromatic heterocyclic residue optionally having a substituent" at Y,

which is preferably pyridyl. The heterocyclic residue may be substituted. As the substituent, exemplified are lower alkyl (as defined for lower alkyl at R¹, R³ and R⁵), lower alkoxy (as defined for lower alkoxy at R³), hydroxy, carboxy, alkoxycarbonyl, halogen atom (chlorine atom, bromine atom, iodine atom and fluorine atom), nitro, amino and the like. When the heterocyclic moiety is substituted, the number of the substituent is preferably 1 or 2.

[0022] The heterocyclic compound [I] and a pharmaceutically acceptable salt thereof are preferably exemplified by the following.

[0023] A heterocyclic compound of the above-mentioned formula [I], wherein

(1) R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent or -COR⁴ wherein R⁴ is hydrogen atom, alkyl optionally having a substituent, aryl optionally having a substituent or arylalkyl optionally having a substituent,

R³ is hydrogen atom, lower alkyl or lower alkoxy,

A is a single bond or >N-R⁵ wherein R⁵ is hydrogen atom or lower alkyl,

B is lower alkylene, and

Y is aryl optionally having a substituent or an aromatic heterocyclic residue optionally having a substituent,

(2) R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl, cycloalkylalkyl, arylalkyl optionally having a substituent, alkenyl, alkynyl, heterocyclic alkyl or

-COR⁴ wherein R⁴ is alkyl, alkenyl or aryl,

R³ is hydrogen atom or lower alkoxy,

A is a single bond or >N-R⁵ wherein R⁵ is lower alkyl,

B is lower alkylene, and

Y is aryl or an aromatic heterocyclic residue optionally having a substituent, or

(3) R¹ is hydrogen atom or lower alkyl,

R² is hydrogen atom, alkyl, cycloalkylalkyl, arylalkyl optionally having a substituent or -COR⁴ wherein R⁴ is alkyl or aryl,


R³ is hydrogen atom,

A is a single bond or >N-R⁵ wherein R⁵ is lower alkyl,


B is lower alkylene, and

Y is an aromatic heterocyclic residue optionally having a substituent and a pharmaceutically acceptable salt thereof.

[0024] In the formula [I], Y-A- is preferably




5




10

15




20

25



30

35



40 wherein


R^A is isopropyl or tert-butyl,

R^B is isopropyl or tert-butyl, and







R^C is isopropyl, tert-butyl, phenyl, thiophen-2-yl, 2-methylpropenyl or 3-butenyl; or

45

50



55




or

wherein

R^A is isopropyl or tert-butyl,

R^B is isopropyl or tert-butyl,

R^C is isopropyl, tert-butyl, phenyl, thiophen-2-yl, 2-methylpropenyl, 3-butenyl, cyclopropyl, 1-butenyl or 2,2-dimethylpropyl, with particular preference given to



[0025] Preferable examples of the heterocyclic compound [I] and the pharmaceutically acceptable salt thereof are

- (1) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (2) 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (3) 2-acetyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (4) 2-methyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (5) 2-hexanoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (6) 2-hexyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (7) 2-isobutyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (8) 2-cyclohexylmethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (9) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(3-phenylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (10) 2-benzoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (11) 2-benzyl-7-[2-(N-methyl-N-(pyridin-2-yl)amino)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (12) 2-benzyl-7-[2-(5-ethyl-pyridin-2-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (13) 2-benzyl-7-[2-(indolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (14) ethyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (15) methyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (16) 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (17) ethyl 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (18) 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(19) ethyl 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,

(20) 2-benzyl-7-[2-(6-carboxyindolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid, and

(21) 2-(4-fluorobenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid, and pharmaceutically acceptable salt thereof. Besides the above-mentioned, preferred are

(22) 2-(2,2-dimethylpropionyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(23) 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(24) 2-benzyl-7-[2-(5-methyl-2-tert-butyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(25) 2-benzyl-7-[2-(5-methyl-2-(thiophen-2-yl)oxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(26) 2-benzyl-7-[2-(5-methyl-2-isopropylloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(27) 2-butyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(28) 2-benzyl-7-[2-[5-methyl-2-(2-methylpropenyl)oxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(29) 2-benzyl-7-[2-[2-(3-butenyl)-5-methyloxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(30) 2-allyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(31) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(2-propynyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(32) 2-(2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(33) 2-benzyl-7-[(indolin-3-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(34) 2-(3-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(35) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-pentanoyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(36) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(4-pentenyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(37) 2-(3-methyl-2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(38) 2-(3,3-dimethylbutyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(39) 2-benzyl-7-methoxy-6-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3RS)-carboxylic acid,

(40) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(pyridin-2-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(41) 2-benzyl-7-(3-methyl-3-phenylbutoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(42) 2-benzyl-7-(3,3-dimethyl-4-phenylbutoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(43) 2-benzyl-7-(2-isopropylbenzoxazol-6-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(44) 2-benzyl-7-(2-tert-butylbenzoxazol-6-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(45) 2-benzyl-7-(2-tert-butylbenzoxazol-5-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(46) 7-(2-tert-butylbenzoxazol-6-yl)methoxy-2-(2,2-dimethylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid, and

(47) 2-benzyl-7-(2-isopropylbenzoxazol-5-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid, and pharmaceutically acceptable salts thereof.

In addition to the above-mentioned, the following heterocyclic compound [I] and pharmaceutically acceptable salts thereof are preferable.

(48) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(pyridin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(49) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-[(pyridin-2-yl)carbonyl]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(50) methyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,


(51) 2-benzyl-7-[2-(2-cyclopropyl-5-methyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(52) 2-(3-methyl-2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

(53) 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-tert-butyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

- (54) 2-benzyl-7-[2-(1-butenyl)-5-methyloxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (55) 2-benzyl-7-[2-(2,2-dimethylpropyl)-5-methyloxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (56) ethyl 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
 (57) 7-(benzofran-2-ylmethoxy)-2-benzyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (58) 2-isobutyryl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (59) 7-[2-(benzofran-2-yl)ethoxy]-2-benzyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (60) 7-[2-(5-ethylpyridin-2-yl)ethoxy]-2-hexanoyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (61) 2-carboxymethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (62) 2-[3-(methoxycarbonyl)propionyl]-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (63) 2-[3-(ethoxycarbonyl)propyl]-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (64) 2-benzyl-6-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3RS)-carboxylic acid,
 (65) 2-(3-acetylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (66) 2-(2-acetylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (67) 2-benzyl-7-[(5-methyl-2-phenyloxazol-4-yl)methoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid.

[0026] Since the heterocyclic compound [I] has an asymmetric carbon at the 3-position of the 1,2,3,4-tetrahydroisoquinoline ring, it includes various stereoisomers. The most preferable configuration is



wherein R¹, R², R³, Y, A and B are as defined above.

[0027] The heterocyclic compound [I] can be converted to a pharmaceutically acceptable salt as necessary.


[0028] When the heterocyclic compound [I] has a basic group, an acid addition salt can be formed. The acid used for forming the acid addition salt is not subject to any particular limitation as long as it can form a salt with a basic moiety and it is a pharmaceutically acceptable acid. Examples of the acid include inorganic acids such as hydrochloric acid, sulfuric acid, phosphoric acid, nitric acid and the like, and organic acids such as oxalic acid, fumaric acid, maleic acid, citric acid, tartaric acid, methanesulfonic acid, p-toluenesulfonic acid and the like, and the like.

[0029] When the heterocyclic compound [I] has an acidic group (e.g., carboxyl group and the like), a salt such as an alkali metal salt (e.g., sodium salt, potassium salt and the like), an alkaline earth metal salt (e.g., calcium salt, magnesium salt and the like) or an organic base salt (e.g., triethylamine salt, dicyclohexylamine salt, pyridine salt and the like) and the like may be formed.

[0030] The heterocyclic compound [I] and the pharmaceutically acceptable salt thereof can be produced according to any of the following Production Methods.

Production Method 1

[0031]



wherein R^1 , R^3 , A, B and Y are as defined above, R^6 is hydrogen atom, alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent, alkenyl optionally having a substituent, alkynyl optionally having a substituent, heterocyclic alkyl optionally having a substituent, $-COR^4$ (R^4 is as defined above) or amino-protecting group, and X is hydroxy, halogen atom (fluorine atom, chlorine atom, bromine atom or iodine atom), or a leaving group such as alkanesulfonyloxy (e.g., methanesulfonyloxy, ethanesulfonyloxy, propanesulfonyloxy, trifluoromethanesulfonyloxy and the like), arylsulfonyloxy (e.g., phenylsulfonyloxy, tolylsulfonyloxy and the like) and the like.

[0032] In Production Method 1, the compound of formula [Ia] (also referred to as compound [Ia]) is synthesized by the reaction of the compound of the formula [II] (also referred to as compound [II]) with the compound of the formula [III] (also referred to as compound [III]).

[0033] The "alkyl optionally having a substituent", "cycloalkyl", "cycloalkylalkyl", "aryl optionally having a substituent", "arylalkyl optionally having a substituent", "alkenyl optionally having a substituent", "alkynyl optionally having a substituent" and "heterocyclic alkyl optionally having a substituent" at R^6 are as defined for R^2 .

[0034] Examples of the amino-protecting group at R^6 are formyl, monochloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, p-nitrobenzyloxycarbonyl, diphenylmethyloxycarbonyl, methoxymethylcarbonyl, methoxymethylloxycarbonyl, trimethylsilyl, 2,2,2-trichloroethoxycarbonyl, 2-methylsulfonylethylloxycarbonyl, tert-butoxycarbonyl (hereinafter to be referred to as Boc), trityl and the like.

Production Method 1-a:

[0035] When X is hydroxy, Production Method 1 comprises a dehydrating reaction such as Mitsunobu Reaction (Reagents for Organic Synthesis, Fisher & Fisher, Vol. 6, 645) and the like. The reaction is generally carried out in a solvent with an azo compound and phosphine. As the azo compound, exemplified are di(C_1 - C_4 alkyl) azodicarboxylate (e.g., diethyl azodicarboxylate and the like), azodicarboxamide (e.g., 1,1'-(azodicarbonyl)dipiperidine and the like) and the like. As the phosphine, exemplified are triarylphosphine (e.g., triphenylphosphine and the like), tri(C_1 - C_4 alkyl) phosphine (e.g., tributylphosphine and the like), and the like.

[0036] As the solvent used in the Production Method 1-a, exemplified are dioxane, acetonitrile, tetrahydrofuran, chloroform, methylene chloride, ethylene chloride, benzene, toluene, xylene, ethyl acetate, N,N-dimethylformamide, N,N-dimethylacetamide, dimethyl sulfoxide and the like; and a mixture thereof and the like, but the solvent is not limited as far as it does not adversely influence the reaction.

[0037] The amount of the compound [II] used in the Production Method 1-a is not subject to any particular limitation. It is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [III]. The azo compound and phosphine are respectively used in an amount of generally 1-3 mol, preferably 1-1.5 mol, per 1 mol of compound [III].

[0038] While the reaction conditions such as reaction temperature and reaction time and the like in Production Method 1-a vary depending on the reagent, solvent and the like to be used in this reaction, the reaction generally proceeds at a temperature of from -30°C to 50°C for 30 min to about a dozen hours.

5 Production Method 1-b:

[0039] When X is halogen atom or leaving group such as alkanesulfonyloxy (e.g., methanesulfonyloxy, ethanesulfonyloxy, propanesulfonyloxy, trifluoromethanesulfonyloxy and the like), arylsulfonyloxy (e.g., phenylsulfonyloxy, tolylsulfonyloxy and the like) and the like, Production Method 1-b is performed in a solvent similar to that used in the
10 Production Method 1-a and in the presence of a base.

[0040] The base used in the Production Method 1-b is exemplified by, but not particularly limited to, inorganic base such as alkali metal carbonate (e.g., sodium carbonate, potassium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate and the like), alkali metal hydroxide (e.g., sodium hydroxide, potassium hydroxide and the like), metal hydride compound (e.g., sodium hydride, potassium hydride, calcium hydride and the like) and the like; organic
15 base such as alkali metal alkoxide (e.g., sodium methoxide, sodium ethoxide, potassium tert-butoxide and the like), amines (e.g., triethylamine, diisopropylethylamine and the like) and the like.

[0041] The amount of the compound [II] used in the Production Method 1-b is not subject to any particular limitation. It is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [III]. The base is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [III].


[0042] While the reaction conditions such as reaction temperature and reaction time and the like in Production Method 1-b vary depending on the reagent, solvent and the like to be used in this reaction, the reaction generally proceeds at a temperature of from -30°C to 150°C for 30 min to about a dozen hours.

[0043] In the Production Method 1, R¹ of compound [III] is preferably lower alkyl. In this case, compound [Ia] wherein R¹ is lower alkyl is obtained, and this compound may be led to the compound [Ia] wherein R¹ is hydrogen atom, by
25 hydrolysis according to a method known per se.

[0044] In the Production Method 1, when R⁶ of compound [III] is amino-protecting group, compound [Ia] wherein R⁶ is amino-protecting group is obtained. This compound may be led to the compound [Ia] wherein R⁶ is hydrogen atom, by deprotection according to a method known per se.

Production Method 2

[0045]



wherein R^1 , R^3 , A, B and Y are as defined above, R^7 is amino-protecting group, R^8 is alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent, alkenyl optionally having a substituent, alkynyl optionally having a substituent or heterocyclic alkyl optionally having a substituent, and U is halogen atom (fluorine atom, chlorine atom, bromine atom, iodine atom), or a leaving group such as alkanesulfonyloxy (e.g., methanesulfonyloxy, ethanesulfonyloxy, propanesulfonyloxy, trifluoromethanesulfonyloxy and the like), arylsulfonyloxy (e.g., phenylsulfonyloxy, tolylsulfonyloxy and the like) and the like.

[0046] In Production Method 2, the amino-protecting group R^7 of a compound of the formula [Ib] (also referred to as compound [Ib]) is eliminated by a method known *per se* to give a compound of the formula [Ic] (also referred to as compound [Ic]), which is then reacted with a compound of the formula [V] (also referred to as compound [V]), whereby a compound of the formula [Id] (also referred to as compound [Id]) is produced.

[0047] The "alkyl optionally having a substituent", "cycloalkyl", "cycloalkylalkyl", "aryl optionally having a substituent", "arylalkyl optionally having a substituent", "alkenyl optionally having a substituent", "alkynyl optionally having a substituent" and "heterocyclic alkyl optionally having a substituent" at R^8 are as defined for R^2 .

[0048] The amino-protecting group at R^7 is as defined for R^6 .

[0049] In the Production Method 2, compound [Ic] is reacted with compound [V] in a solvent that does not inhibit the reaction, such as dioxane, acetonitrile, tetrahydrofuran, chloroform, methylene chloride, ethylene chloride, benzene, toluene, xylene, ethyl acetate, N,N-dimethylformamide, N,N-dimethylacetamide, dimethyl sulfoxide and the like; and a mixture thereof and the like, in the presence of a base, to give compound [Id].

[0050] In the Production Method 2, the base used for the reaction of compound [Ic] with compound [V] is exemplified, but not particularly limited to, inorganic base such as alkali metal carbonate (e.g., sodium carbonate, potassium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate and the like), alkali metal hydroxide (e.g., sodium hydroxide, potassium hydroxide and the like), metal hydride compound (e.g., sodium hydride, potassium hydride and calcium hydride and the like) and the like; and organic base such as alkali metal alkoxide (e.g., sodium methoxide,

sodium ethoxide, potassium tert-butoxide and the like), amines (e.g., triethylamine, diisopropylethylamine and the like) and the like.


[0051] The amount of the compound [V] used in the Production Method 2 is generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [Ic]. The base is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [Ic].

[0052] While the reaction conditions such as reaction temperature and reaction time and the like for the reaction of compound [Ic] with compound [V] in the Production Method 2 vary depending on the reagent, solvent and the like to be used in this reaction, the reaction generally proceeds at a temperature of from -30°C to 150°C for 30 min to about two dozen hours.

[0053] In the Production Method 2, R¹ of compound [Ib] is preferably lower alkyl. In this case, compound [Id] wherein R¹ is lower alkyl is obtained, and this compound may be led to the compound [Id] wherein R¹ is hydrogen atom, by hydrolysis according to a method known *per se*.

Production Method 3

[0054]



wherein R¹, R³, R⁴, A, B and Y are as defined above.

[0055] In Production Method 3, a compound of the formula [Ie] (also referred to as compound [Ie]) is synthesized by reacting compound [Ic] with a compound of the formula [VI] (also referred to as compound [VI]).

[0056] In the Production Method 3, compound [VI] may be used not only in the form of a free acid, but also in the form of a salt (e.g., sodium salt, potassium salt, calcium salt, triethylamine salt, pyridine salt and the like) or as a reactive derivative (e.g., acid halide such as acid chloride, acid bromide and the like; acid anhydride; mixed acid anhydride with substituted phosphoric acid such as dialkyl phosphate and the like, and alkyl carbonate such as monoethyl carbonate and the like; reactive amide which is amide with imidazole and the like; ester such as cyanomethyl ester and 4-nitro-

phenyl ester and the like), and the like.

[0057] In the Production Method 3, when the compound [VI] is used in the form of a free acid or a salt, this reaction is preferably carried out in the presence of a condensing agent. As the condensing agent, a dehydrating agent such as carbodiimide compound (e.g., N,N'-dicyclohexylcarbodiimide, 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide, N-cyclohexyl-N'-morpholinoethylcarbodiimide, N-cyclohexyl-N'-(4-diethylaminocyclohexyl)carbodiimide and the like), azolide compound (e.g., N,N'-carbonyldiimidazole, N,N'-thionylidiimidazole and the like) and the like can be used. The condensing agent is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [Ic]. It is considered that, when a condensing agent is used, compound [VI] becomes a reactive derivative and proceeds to the reaction.

[0058] The Production Method 3 is generally performed in a solvent inert to the reaction. As the solvent, exemplified are acetone, dioxane, acetonitrile, chloroform, benzene, methylene chloride, ethylene chloride, tetrahydrofuran, ethyl acetate, N,N-dimethylformamide, pyridine, water and a mixed solvent thereof. In the Production Method 3, a base such as triethylamine, pyridine, 4-dimethylaminopyridine, potassium carbonate and the like can be used. When the base is used, it is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [Ic].

[0059] In Production Method 3, compound [VI] is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [Ic].

[0060] While the reaction conditions such as reaction temperature and reaction time and the like in the reaction of compound [VI] with compound [Ic] in the Production Method 3 vary depending on the reagent, solvent and the like to be used in this reaction, the reaction generally proceeds at a temperature of from -30°C to 150°C for 10 min to about a dozen hours.


[0061] In the Production Method 3, compound [Ie'] is obtained by reacting compound [VI] with compound [Ic], isolating and reducing the resulting compound [Ie]. This reduction reaction is carried out in a solvent that does not adversely influence the reaction (e.g., water, methanol, ethanol, dioxane, acetonitrile, tetrahydrofuran, chloroform, methylene chloride, ethylene chloride, benzene, toluene, xylene, ethyl acetate, N,N-dimethylformamide, N,N-dimethylacetamide, dimethyl sulfoxide, a mixture thereof and the like) in the presence of a reducing agent.

[0062] The reducing agent used in this reduction is exemplified by metal hydride complex (e.g., lithium aluminum hydride, sodium cyanoborohydride (NaBH_3CN), sodium borohydride and the like), borane and the like. The agent is not subject to any particular limitation as long as it is generally used for reducing a carbonyl group to a methylene group. The reducing agent is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [Ie].

[0063] While the reaction conditions such as reaction temperature and reaction time and the like in the reduction reaction vary depending on the reagent, solvent and the like to be used in this reaction, the reaction generally proceeds at a temperature of from -30°C to 150°C for 30 min to about a dozen hours.

Production Method 4

[0064]



wherein R^1 , R^2 , R^3 , R^5 , B, X, Y and U are as defined above, and R^{10} is an amino-protecting group.

[0065] The amino-protecting group at R^{10} is as defined for R^6 .

[0066] Production Method 4 is intended for the heterocyclic compound [I] wherein A is $>N-R^5$. In this method, compound of the formula [VII] (also referred to as compound [VII]) is reacted with compound of the formula [VIII] (also referred to as compound [VIII]) in the same manner as in Production Method 1 to give a compound of the formula [IX] (also referred to as compound [IX]), and the amino-protecting group at R^{10} of the compound [IX] is eliminated by a

method known *per se* to give a compound of the formula [X] (also referred to as compound [X]), which is then reacted with a compound of the formula [XI] (also referred to as compound [XI]), whereby a compound of the formula [If] (also referred to as compound [If]) is produced.

[0067] The reaction of compound [X] with compound [XI] in Production Method 4 is carried out in a solvent that does not adversely influence the reaction in the presence of a base. As the solvent, exemplified are dioxane, acetonitrile, tetrahydrofuran, chloroform, methylene chloride, ethylene chloride, benzene, toluene, xylene, ethyl acetate, N,N-dimethylformamide, N,N-dimethylacetamide, dimethyl sulfoxide and the like; and a mixture thereof. When compound [XI] can be used as a solvent, it can be used as a solvent.


[0068] The base used in the reaction of compound [X] with compound [XI] in the Production Method 4 is exemplified, but not particularly limited to, inorganic base such as alkali metal carbonate (e.g., sodium carbonate, potassium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate and the like), alkali metal hydroxide (e.g., sodium hydroxide, potassium hydroxide and the like), metal hydride compound (e.g., sodium hydride, potassium hydride, calcium hydride and the like) and the like; and organic base such as alkali metal alkoxide (e.g., sodium methoxide, sodium ethoxide, potassium tert-butoxide and the like) and amines (e.g., triethylamine, diisopropylethylamine and the like) and the like. The base is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [X].

[0069] In Production Method 4, compound [XI] is used in a proportion of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [X].

[0070] While the reaction conditions such as reaction temperature and reaction time and the like in the reaction of compound [X] with compound [XI] in the Production Method 4 vary depending on the reagent, solvent and the like to be used in this reaction, the reaction generally proceeds at a temperature of from -30°C to 150°C for 30 min to about a dozen hours.

Production Method 5

[0071]



wherein R^1 , R^3 , A, B and Y are as defined above, R^9 is such a group as makes a group of the formula: $-CH_2-R^9$ alkyl optionally having a substituent, cycloalkylalkyl, arylalkyl optionally having a substituent, alkenyl optionally having a substituent, alkynyl optionally having a substituent or heterocyclic alkyl optionally having a substituent.

[0072] In Production Method 5, compound [Ic] is reacted with a compound of the formula [XII] (also referred to as compound [XII]), whereby a compound of the formula [Ig] (also referred to as compound [Ig]) is produced.

[0073] In a group of the formula: $-CH_2-R^9$, the "alkyl optionally having a substituent", "cycloalkylalkyl", "arylalkyl optionally having a substituent", "alkenyl optionally having a substituent", "alkynyl optionally having a substituent" and "heterocyclic alkyl optionally having a substituent" are as defined for R^2 .

[0074] In Production Method 5, compound [Ic] and compound [XII] are condensed in a solvent which does not inhibit the reaction (e.g., water, methanol, ethanol, dioxane, acetonitrile, tetrahydrofuran, chloroform, methylene chloride, ethylene chloride, benzene, toluene, xylene, ethyl acetate, N,N-dimethylformamide, N,N-dimethylacetamide, dimethyl sulfoxide, a mixture thereof and the like) in the presence of a reducing agent.

[0075] The reducing agent used in the Production Method 5 is not subject to any particular limitation, and is exemplified by metal hydride complex (e.g., lithium aluminum hydride, sodium cyanoborohydride (NaBH_3CN), sodium borohydride and the like), borane and the like.

[0076] In Production Method 5, compound [XII] is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [Ic]. The reducing agent is used in an amount of generally 1-5 mol, preferably 1-3 mol, per 1 mol of compound [Ic].

[0077] While the reaction conditions such as reaction temperature and reaction time and the like in Production Method 5 vary with the reagent, solvent and the like to be used in this reaction, the reaction generally proceeds at a temperature of from -30°C to 150°C for 30 min to about a dozen hours.

[0078] The heterocyclic compound [I] obtained in the above-mentioned Production Methods 1-5 may be isolated by a conventional method, and, where necessary, purified by a conventional method such as recrystallization, preparative thin-layer chromatography, column chromatography and the like. The compound can be also purified into a salt as necessary.

[0079] The heterocyclic compound [I] may be converted to a pharmaceutically acceptable salt thereof by a method known *per se*.

[0080] The pharmaceutical composition comprising the heterocyclic compound [I] or a pharmaceutically acceptable salt thereof of the present invention may contain an additive and the like. As an additive, exemplified are excipient (e.g., starch, lactose, sugar, calcium carbonate, calcium phosphate and the like), binder (e.g., starch, gum arabic, carboxymethylcellulose, hydroxypropylcellulose, crystalline cellulose and the like), lubricant (e.g., magnesium stearate, talc and the like), disintegrator (e.g., carboxymethylcellulose calcium, talc and the like), and the like.

[0081] The above-mentioned components are mixed to give a preparation for oral administration such as capsule, tablet, powder, granule, dry syrup and the like, or a preparation for parenteral administration such as injection, suppository and the like, according to a method known *per se*.

[0082] While the dose of the heterocyclic compound [I] or a pharmaceutically acceptable salt thereof may vary according to the administration subject, symptom and other factors, when it is orally administered to an adult patient with, for example, diabetes, diabetic complications or hyperlipidemia, the single dose is approximately 1 - 500 mg/kg body weight, which is administered 1 to 3 times a day.

[0083] The heterocyclic compound [I] and a pharmaceutically acceptable salt of the present invention show a superior hypoglycemic action, a blood hypolipidemic action, an insulin resistance-improving action and a PPAR activating action in mammals (human, cattle, horse, dog, cat, rat, mouse, hamster and the like), and are useful as a hypoglycemic agent, a hypolipidemic agent, an insulin resistance improver, a therapeutic agent of diabetes, a therapeutic agent of diabetic complication, a glucose tolerance improver, an anti-atherosclerosis agent, an anti-obesity agent, an antiinflammatory agent, an agent for the prophylaxis or treatment of PPAR-mediated disease and an agent for the prophylaxis or treatment of syndrome X. To be specific, the heterocyclic compound [I] and a pharmaceutically acceptable salt of the present invention are useful for the prophylaxis or treatment of diabetes, diabetic complications, hyperlipidemia, atherosclerosis, hyperglycemia, diseases caused by insulin resistant impaired glucose tolerance, diseases caused by insulin resistance, obesity, inflammation, PPAR-mediated disease and syndrome X.

Examples

[0084] The present invention is explained in detail by referring to the following examples. The present invention is not limited by these examples in any way.

Example 1

Sodium 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0085]

(1) Ethyl 2-tert-butoxycarbonyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.50 g) and 2-(5-methyl-2-phenyloxazol-4-yl)ethyl methanesulfonate (2.50 g) were dissolved in N,N-dimethylformamide (20 ml). There-to was added potassium carbonate (2.0 g) and the mixture was stirred at 80°C for 5 hr. Ethyl acetate (100 ml) was added to the reaction mixture, and the mixture was washed with water (100 ml) and saturated brine (50 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-tert-butoxycarbonyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.62 g).

Ethyl 2-tert-butoxycarbonyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v(neat) cm^{-1} ; 2978, 2930, 1738, 1699, 1614, 1587.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.29 (3H, t, $J=7.0\text{Hz}$), 1.46, 1.50 (9H, s, s), 2.36 (3H, s), 2.95 (2H, t, $J=6.8\text{Hz}$), 2.90-3.30 (2H, m), 4.00-4.40 (4H, m), 4.51, 4.61 (2H, s, s), 4.70-4.90, 5.00-5.20 (1H, m, m), 6.60-6.90 (2H, m), 7.12 (1H, d, $J=8.4\text{Hz}$), 7.30-7.55 (3H, m), 7.90-8.15 (2H, m).

(2) The compound (5.2 g) obtained in (1) above was dissolved in formic acid (20 ml). Thereto was added a 8.78N hydrogen chloride solution (6.0 ml) in 2-propanol under ice-cooling, and the mixture was stirred at room temperature for 10 min. Ethyl acetate (100 ml) was added to the reaction mixture, and the mixture was neutralized with saturated aqueous sodium hydrogencarbonate solution and separated into two layers. The obtained ethyl acetate layer was washed with saturated brine (50 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure to give ethyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (3.6 g). Ethyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v (nujol) cm^{-1} ; 3476, 1742, 1639, 1611, 1553.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.29 (3H, t, $J=7.0\text{Hz}$), 2.02 (1H, s), 2.36 (3H, s), 2.80-3.10 (4H, m), 3.50-3.80 (1H, m), 4.00-4.40 (6H, m), 6.50-6.80 (2H, m), 7.00 (1H, d, $J=8.4\text{Hz}$), 7.30-7.50 (3H, m), 7.90-8.10 (2H, m).

(3) The compound (1.11 g) obtained in (2) above was dissolved in methanol (20 ml). Thereto was added 1N aqueous sodium hydroxide solution (3.0 ml) and the mixture was stirred at room temperature for 2 hr. The solvent was evaporated under reduced pressure. Water (5 ml) was added to the obtained crystalline residue, and the crystals were collected by filtration to give the title compound (0.92 g).

IR v (nujol) cm^{-1} ; 3427, 1589, 1504.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

2.35 (3H, s), 2.60-3.10 (6H, m), 3.86 (2H, br-s), 4.14 (2H, t, $J=6.6\text{Hz}$), 6.50-6.80 (2H, m), 6.94 (1H, d, $J=8.1\text{Hz}$), 7.40-7.60 (3H, m), 7.75-8.05 (2H, m).

Example 2

2-Benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0086]

(1) The compound (1.40 g) obtained in Example 1(2) was dissolved in N,N-dimethylformamide (20 ml). Thereto was added sodium hydride (160 mg, 60% suspension in oil) under ice-cooling, and the mixture was stirred at room temperature for 20 min. To the obtained mixture was dropwise added benzyl bromide (0.40 ml), and the mixture was stirred further at the same temperature for 1 hr. Ethyl acetate (50 ml) was added to the reaction mixture, and the mixture was washed with water (50 ml) and saturated brine (30 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.38 g).

Ethyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v (nujol) cm^{-1} ; 1728, 1639, 1614, 1551.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.21 (3H, t, $J=7.0\text{Hz}$), 2.34 (3H, s), 2.92 (2H, t, $J=7.0\text{Hz}$), 3.05-3.20 (2H, m), 3.60-4.00 (5H, m), 4.12 (2H, q, $J=7.0\text{Hz}$), 4.16 (2H, t, $J=7.0\text{Hz}$), 6.51 (1H, d, $J=2.0\text{Hz}$), 6.68 (1H, dd, $J=2.0, 8.4\text{Hz}$), 6.99 (1H, d, $J=8.4\text{Hz}$), 7.30-7.50 (8H, m), 7.80-8.10 (2H, m).

(2) The compound (8.20 g) obtained in (1) above was dissolved in 80 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 2N aqueous lithium hydroxide solution (42 ml), and the mixture was stirred at 50°C for 2 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration. The obtained crude crystals (9.0 g) were recrystallized from methanol to give the title compound (6.33 g).

IR v (nujol) cm^{-1} ; 1638, 1501.

¹H-NMR (DMSO-d₆) δ (ppm) ;
 2.33 (3H, s), 2.65-3.30 (4H, m), 3.50-4.00 (5H, m), 4.00-6.20 (1H, br), 4.13 (2H, t, J=7.0Hz), 6.59 (1H, br-s),
 6.68 (1H, br-d, J=8.4Hz), 7.01 (1H, d, J=8.4Hz), 7.32 (5H, s), 7.35-7.70 (3H, m), 7.85-8.10 (2H, m).

Example 3

2-Acetyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0087]

(1) The compound (800 mg) obtained in Example 1(2) was dissolved in methylene chloride (8.0 ml). Thereto was added acetic anhydride (0.23 ml) under ice-cooling, and the mixture was stirred at room temperature for 10 min. Ethyl acetate (30 ml) was added to the reaction mixture, and the mixture was neutralized with saturated aqueous sodium hydrogen carbonate solution, and then the two layers were separated. The obtained organic layer was washed with saturated brine (10 ml) and dried over Na₂SO₄. The solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-acetyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (873 mg).

Ethyl 2-acetyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v (nujol) cm⁻¹ ; 1732, 1651, 1555.

¹H-NMR (CDCl₃) δ (ppm);

1.12 (3H, t, J=7.0Hz), 2.13, 2.21 (3H, s, s), 2.36 (3H, s), 2.95 (2H, t, J=6.6Hz), 3.05-3.30 (2H, m), 4.04 (2H, q, J=7.0Hz), 4.22 (2H, t, J=6.6Hz), 4.62 (2H, s), 5.45 (1H, dd, J=4.0, 5.7Hz), 6.55-6.85 (2H, m), 7.04 (1H, d, J=8.2Hz), 7.30-7.50 (3H, m), 7.85-8.10 (2H, m).

(2) The compound (800 mg) obtained in (1) above was dissolved in 5.0 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (3.0 ml), and the mixture was stirred at room temperature for 30 min. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (668 mg).

IR v (nujol) cm⁻¹ ; 3400, 1732, 1641, 1612, 1555.

¹H-NMR (CDCl₃) δ (ppm) ;

2.10, 2.17 (3H, s, s), 2.32 (3H, s), 2.70-3.30 (4H, m), 3.90-4.20 (2H, m), 4.30-4.90 (2H, m), 5.35-5.60 (1H, m), 6.50-6.80 (2H, m), 7.03 (1H, d, J=8.1Hz), 7.30-7.60 (3H, m), 7.80-8.10 (2H, m).

Example 4

2-Methyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0088]

(1) The compound (1.0 g) obtained in Example 1(2) was dissolved in methanol (10 ml). Thereto were added formalin (0.4 ml) and sodium cyanoborohydride (310 mg), and the mixture was stirred at room temperature for 1 hr. Methanol was evaporated under reduced pressure. Ethyl acetate (20 ml) was added to the obtained residue, and the mixture was washed with water (20 ml) and saturated brine (10 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-methyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.08 g).

Ethyl 2-methyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v(neat) cm⁻¹ ; 2926, 2874, 1732, 1641, 1614.

¹H-NMR (CDCl₃) δ (ppm);

1.25 (3H, t, J=7.0Hz), 2.35 (3H, s), 2.50 (3H, s), 2.94 (2H, t, J=6.8Hz), 3.02 (2H, d, J=6.0Hz), 3.45 (1H, t, J=6.0Hz), 3.64 (1H, d, J=15.6Hz), 3.98 (1H, d, J=15.6Hz), 4.17 (2H, q, J=7.0Hz), 4.20 (2H, t, J=6.6Hz), 6.56 (1H, d, J=2.0Hz), 6.70 (1H, dd, J=2.0, 8.4Hz), 6.98 (1H, d, J=8.4Hz), 7.30-7.50 (3H, m), 7.85-8.10 (2H, m).

(2) The compound (1.08 g) obtained in (1) above was dissolved in 10 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (7.5 ml), and the mixture was stirred at

room temperature for 1 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (0.74 g).

IR ν (nujol) cm^{-1} ; 1616, 1555, 1541, 1506.

$^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ (ppm);

2.36 (3H, s), 2.55 (3H, s), 2.70-3.10 (4H, m), 3.40-3.60 (1H, m), 3.70-4.30 (4H, m), 6.60-6.80 (2H, m), 7.05 (1H, d, $J=8.6\text{Hz}$), 7.35-7.65 (3H, m), 7.75-8.10 (2H, m).

Example 5

2-Hexanoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0089]

(1) The compound (1.0 g) obtained in Example 1(2) was dissolved in methylene chloride (10 ml). Thereto were added hexanoyl chloride (0.41 ml) and triethylamine (0.51 ml) under ice-cooling, and the mixture was stirred at the same temperature for 10 min. Ethyl acetate (70 ml) was added to the reaction mixture, and the mixture was washed with 10% aqueous citric acid solution (50 ml), saturated aqueous sodium hydrogencarbonate solution (50 ml) and then saturated brine (50 ml) and dried over Na_2SO_4 . The solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-hexanoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.02 g).

Ethyl 2-hexanoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (nujol) cm^{-1} ; 1736, 1653, 1587.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

0.70-1.90 (12H, m), 2.20-2.60 (2H, m), 2.36 (3H, s), 2.95 (2H, t, $J=6.8\text{Hz}$), 3.10-3.20 (2H, m), 4.04 (2H, q, $J=7.0\text{Hz}$), 4.22 (2H, t, $J=7.0\text{Hz}$), 4.63 (2H, s), 5.45 (1H, dd, $J=4.0, 5.4\text{Hz}$), 6.60-6.90 (2H, m), 7.04 (1H, d, $J=8.1\text{Hz}$), 7.30-7.50 (3H, m), 7.80-8.10 (2H, m).

(2) The compound (1.02 g) obtained in (1) above was dissolved in 10 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (6.0 ml), and the mixture was stirred at room temperature for 4 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (0.56 g).

IR ν (nujol) cm^{-1} ; 1742, 1641, 1612, 1572.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

0.88 (3H, br-t, $J=6.0\text{Hz}$), 1.10-1.90 (6H, m), 2.32 (3H, s), 2.30-2.50 (2H, m), 2.70-3.30 (4H, m), 4.07 (2H, t, $J=7.0\text{Hz}$), 4.60 (2H, s), 5.40-5.60 (1H, m), 6.60-6.80 (2H, m), 7.05 (1H, d, $J=8.6\text{Hz}$), 7.35-7.65 (3H, m), 7.75-8.10 (2H, m).

Example 6

2-Hexyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0090]

(1) The compound (1.20 g) obtained in Example 1(2) was dissolved in N,N-dimethylformamide (12 ml). Thereto were added hexyl iodide (0.65 ml) and potassium carbonate (0.82 g) and the mixture was stirred at 50°C for 15 hr. Water (100 ml) was added to the reaction mixture, and the mixture was extracted twice with ethyl acetate (50 ml). The ethyl acetate layer was washed with saturated brine (100 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-hexyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.05 g).

Ethyl 2-hexyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (nujol) cm^{-1} ; 1724, 1643, 1611.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

0.88 (3H, br-t, $J=5.0\text{Hz}$), 1.10-1.75 (11H, m), 2.35 (3H, s), 2.50-2.80 (2H, m), 2.94 (2H, t, $J=7.0\text{Hz}$), 3.02 (2H,

d, J=5.5Hz), 3.68 (1H, t, J=5.5Hz), 3.83 (1H, s), 3.94 (1H, s), 4.12 (2H, q, J=7.0Hz), 4.20 (2H, t, J=7.0Hz), 6.50-6.80 (2H, m), 6.97 (1H, d, J=8.4Hz), 7.30-7.60 (3H, m), 7.80-8.10 (2H, m).

(2) The compound (1.0 g) obtained in (1) above was dissolved in 10 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 2N aqueous lithium hydroxide solution (5.1 ml), and the mixture was stirred at room temperature for 11 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (0.80 g).

IR v (nujol) cm^{-1} ; 1620, 1555, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

0.89 (3H, br-t, J=6.0Hz), 1.00-1.45 (6H, m), 1.50-1.90 (2H, m), 2.36 (3H, s), 2.70-3.30 (2H, m), 2.93 (2H, t, J=6.2Hz), 3.15 (2H, d, J=6.4Hz), 3.75 (1H, t, J=6.4Hz), 4.00-4.40 (4H, m), 6.23 (1H, br-s), 6.60-6.85 (2H, m), 7.06 (1H, d, J=8.4Hz), 7.30-7.60 (3H, m), 7.80-8.10 (2H, m).

Example 7

2-Isobutyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0091]

(1) The compound (1.40 g) obtained in Example 1(2) was dissolved in N,N-dimethylformamide (14 ml). Thereto were added isobutyl iodide (1.20 ml) and potassium carbonate (0.95 g), and the mixture was stirred at 50°C for 3 days. Water (100 ml) was added to the reaction mixture, and the mixture was extracted twice with ethyl acetate (50 ml). The obtained ethyl acetate layer was washed with saturated brine (100 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-isobutyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.15 g).

Ethyl 2-isobutyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v (nujol) cm^{-1} ; 1719, 1645, 1614, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

0.89 (6H, d, J=6.6Hz), 1.19 (3H, t, J=7.0Hz), 1.70-1.90 (1H, m), 2.35 (3H, s), 2.10-2.50 (2H, m), 2.94 (2H, t, J=7.0Hz), 3.02 (2H, d, J=5.2Hz), 3.66 (1H, t, J=5.2Hz), 3.83 (1H, s), 3.92 (1H, s), 4.16 (2H, q, J=7.0Hz), 4.20 (2H, t, J=7.0Hz), 6.45-6.75 (2H, m), 6.97 (1H, d, J=8.2Hz), 7.25-7.50 (3H, m), 7.85-8.10 (2H, m).

(2) The compound (1.05 g) obtained in (1) above was dissolved in 10 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 2N aqueous lithium hydroxide solution (5.7 ml), and the mixture was stirred at room temperature for 24 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (0.83 g).

IR v (nujol) cm^{-1} ; 3339, 1620, 1553, 1508.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

0.95 (3H, d, J=6.7Hz), 1.01 (3H, d, J=7.0Hz), 1.95-2.15 (1H, m), 2.36 (3H, s), 2.60-2.80 (2H, m), 2.94 (2H, t, J=6.7Hz), 3.16 (2H, d, J=6.6Hz), 3.70 (1H, t, J=6.6Hz), 4.11 (2H, s), 4.18 (2H, t, J=6.6Hz), 4.84 (1H, br-s), 6.60-6.90 (2H, m), 7.49 (1H, d, J=8.1Hz), 7.25-7.50 (3H, m), 7.85-8.10 (2H, m).

Example 8

2-Cyclohexylmethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0092]

(1) The compound (1.40 g) obtained in Example 1(2) was dissolved in N,N-dimethylformamide (14 ml). Thereto were added cyclohexylmethyl bromide (1.44 ml) and potassium carbonate (0.95 g), and the mixture was stirred at 50°C for 2 days. Water (100 ml) was added to the reaction mixture, and the mixture was extracted twice with ethyl acetate (50 ml). The ethyl acetate layer was washed with saturated brine (100 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-cyclohexylmethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquino-

line-(3S)-carboxylate (1.0 g).

Ethyl 2-cyclohexylmethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v (nujol) cm^{-1} ; 1728, 1638, 1614, 1504.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

0.80-2.00 (11H, m), 1.19 (3H, t, $J=7.0\text{Hz}$), 2.36 (3H, s), 2.40-2.55 (2H, m), 2.94 (2H, t, $J=7.1\text{Hz}$), 3.00 (2H, d, $J=5.3\text{Hz}$), 3.65 (1H, t, $J=5.3\text{Hz}$), 3.82 (1H, s), 3.92 (1H, s), 4.05 (2H, t, $J=7.0\text{Hz}$), 4.16 (2H, q, $J=7.0\text{Hz}$), 6.50-6.80 (2H, m), 6.97 (1H, d, $J=8.1\text{Hz}$), 7.30-7.50 (3H, m), 7.80-8.10 (2H, m).

(2) The compound (0.95 g) obtained in (1) above was dissolved in 28 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 2N aqueous lithium hydroxide solution (7.13 ml), and the mixture was stirred at room temperature for 3 days. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (0.76 g).

IR v (nujol) cm^{-1} ; 3319, 1624, 15.06.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

0.70-2.10 (11H, m), 2.36 (3H, s), 2.40-2.55 (2H, m), 2.93 (2H, t, $J=6.4\text{Hz}$), 3.16 (2H, d, $J=7.2\text{Hz}$), 3.70 (1H, t, $J=7.2\text{Hz}$), 4.00-4.30 (4H, m), 5.30 (1H, br-s), 6.60-6.90 (2H, m), 7.08 (1H, d, $J=8.4\text{Hz}$), 7.30-7.50 (3H, m), 7.80-8.10 (2H, m).

Example 9

7-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-(3-phenylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0093]

(1) The compound (1.40 g) obtained in Example 1(2) was dissolved in N,N-dimethylformamide (14 ml). Thereto were added 3-phenylpropyl bromide (0.78 ml) and potassium carbonate (0.95 g), and the mixture was stirred at 50°C for 22 hr. Water (100 ml) was added to the reaction mixture, and the mixture was extracted twice with ethyl acetate (50 ml). The ethyl acetate layer was washed with saturated brine (100 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(3-phenylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.05 g).

Ethyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(3-phenylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v (nujol) cm^{-1} ; 1720, 1647, 1612, 1504.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

1.17 (3H, t, $J=7.0\text{Hz}$), 1.60-2.05 (2H, m), 2.35 (3H, s), 2.50-2.80 (4H, m), 2.94 (2H, t, $J=7.1\text{Hz}$), 3.04 (2H, d, $J=5.7\text{Hz}$), 3.67 (1H, t, $J=5.7\text{Hz}$), 3.84 (1H, s), 3.94 (1H, s), 4.04 (2H, t, $J=7.1\text{Hz}$), 4.16 (2H, q, $J=7.0\text{Hz}$), 6.50-6.80 (2H, m), 7.07 (1H, d, $J=9.0\text{Hz}$), 7.20 (5H, s), 7.10-7.50 (3H, m), 7.80-8.10 (2H, m).

(2) The compound (1.0 g) obtained in (1) above was dissolved in 10 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 2N aqueous lithium hydroxide solution (4.77 ml), and the mixture was stirred at room temperature for 10 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (0.66 g).

IR v (nujol) cm^{-1} ; 3346, 1616, 1556, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

1.70-2.20 (3H, m), 2.35 (3H, s), 2.40-2.70 (4H, m), 2.70-3.30 (2H, m), 2.92 (2H, t, $J=6.3\text{Hz}$), 3.10 (2H, d, $J=7.0\text{Hz}$), 3.65 (1H, t, $J=7.0\text{Hz}$), 3.90-4.30 (4H, m), 5.12 (1H, br-s), 6.55-6.80 (2H, m), 6.90-7.25 (6H, m), 7.25-7.55 (3H, m), 7.80-8.10 (2H, m).

Example 10

2-Benzoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0094]

(1) The compound (1.4 g) obtained in Example 1(2) was dissolved in methylene chloride (14 ml). Thereto were added benzoyl chloride (0.48 ml) and triethylamine (0.72 ml) under ice-cooling, and the mixture was stirred at the same temperature for 15 min. Ethyl acetate (100 ml) was added to the reaction mixture, and the mixture was washed successively with 10% aqueous citric acid solution (50 ml), saturated aqueous sodium hydrogencarbonate solution (50 ml) and then saturated brine (50 ml) and dried over Na_2SO_4 . The solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-benzoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.16 g).

Ethyl 2-benzoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v (nujol) cm^{-1} ; 1734, 1638, 1612, 1591.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

0.75-1.15 (3H, m), 2.35 (3H, s), 2.93 (2H, t, $J=6.6\text{Hz}$), 3.05-3.25 (2H, m), 3.85-4.40 (4H, m), 4.20-4.80 (2H, m), 5.00-5.60 (1H, m), 6.47 (1H, br-s), 6.72 (1H, br-d, $J=8.4\text{Hz}$), 7.05 (1H, br-d, $J=8.4\text{Hz}$), 7.30-7.60 (8H, m), 7.80-8.10 (2H, m).

(2) The compound (1.0 g) obtained in (1) above was dissolved in 10 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (6.0 ml), and the mixture was stirred at room temperature for 1.5 hr. The solvent was evaporated under reduced pressure. Thereto was added water (20 ml), and the mixture was washed with ethyl acetate (10 ml). The obtained aqueous layer was acidified with 6N hydrochloric acid, and the mixture was extracted twice with diethyl ether (20 ml). The diethyl ether layer was washed with saturated brine (30 ml) and dried over Na_2SO_4 . Diethyl ether was evaporated under reduced pressure. The obtained residue was recrystallized from methanol to give the title compound (0.75 g).

IR v (nujol) cm^{-1} ; 1730, 1636, 1551.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

2.32 (3H, s), 2.87 (2H, t, $J=6.4\text{Hz}$), 3.00-3.35 (2H, m), 4.02 (2H, t, $J=6.4\text{Hz}$), 4.40-4.90 (2H, m), 4.90-5.30 (1H, br), 5.00-5.65 (1H, m), 6.40 (1H, br-s), 6.50-6.80 (1H, m), 7.03 (1H, d, $J=8.4\text{Hz}$), 7.20-7.60 (8H, m), 7.75-8.05 (2H, m).

Example 11

Sodium 2-benzyl-7-[2-(N-methyl-N-(pyridin-2-yl)amino)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0095]

(1) Ethyl 2-benzyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S) -carboxylate (1.38 g) was dissolved in N,N-dimethylformamide (10 ml). Thereto was added sodium hydride (210 mg, 60% suspension in oil) under ice-cooling and the mixture was stirred at room temperature for 30 min. Thereto was added 2-(N-tert-butoxycarbonyl-N-methylamino)ethyl methanesulfonate (1.30 g) and the mixture was stirred further at the same temperature for 1 hr. Ethyl acetate (50 ml) was added to the reaction mixture, and the mixture was washed with water (50 ml) and saturated brine (30 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-benzyl-7-[2-(N-tert-butoxycarbonyl-N-methylamino)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.44 g).

Ethyl 2-benzyl-7-[2-(N-tert-butoxycarbonyl-N-methylamino)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR v(neat) cm^{-1} ; 2978, 2932, 1732, 1695, 1614.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.23 (3H, t, $J=7.0\text{Hz}$), 1.44 (9H, s), 2.95 (3H, s), 3.08 (2H, d, $J=4.9\text{Hz}$), 3.54 (2H, t, $J=5.5\text{Hz}$), 3.60-4.30 (7H, m), 6.50 (1H, d, $J=2.0\text{Hz}$), 6.68 (1H, dd, $J=2.0, 8.1\text{Hz}$), 7.01 (1H, d, $J=8.1\text{Hz}$), 7.20-7.50 (5H, m).

(2) The compound (1.44 g) obtained in (1) above was dissolved in formic acid (7.0 ml). Thereto was added 8.78N

hydrogen chloride solution (2.0 ml) in 2-propanol, and the mixture was stirred at room temperature for 15 min. Ethyl acetate (100 ml) was added to the reaction mixture, and the mixture was neutralized with saturated aqueous sodium hydrogencarbonate solution, and then the two layers were separated. The obtained ethyl acetate layer was washed with saturated brine (50 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure to give ethyl 2-benzyl-7-(2-methylaminoethoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.08 g).

Ethyl 2-benzyl-7-(2-methylaminoethoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (neat) cm⁻¹; 3332, 1732, 1612, 1504.

¹H-NMR (CDCl₃) δ (ppm);

1.22 (3H, t, J=7.0Hz), 2.41 (1H, br-s), 2.49 (3H, s), 2.95 (2H, t, J=5.5Hz), 3.08 (2H, d, J=4.9Hz), 3.60-4.25 (7H, m), 6.52 (1H, d, J=2.0Hz), 6.70 (1H, dd, J=2.0, 8.4Hz), 7.00 (1H, d, J=8.4Hz), 7.20-7.50 (5H, m).

(3) The compound (1.05 g) obtained in (2) above was dissolved in 2-chloropyridine (2.0 ml), and the mixture was stirred at 140°C for 16 hr. The reaction mixture was purified by silica gel column chromatography to give ethyl 2-benzyl-7-[2-(N-methyl-N-(pyridin-2-yl)amino)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (0.5 g).

Ethyl 2-benzyl-7-[2-(N-methyl-N-(pyridin-2-yl)amino)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (neat) cm⁻¹; 2932, 2905, 1732, 1597, 1560, 1504.

¹H-NMR (CDCl₃) δ (ppm);

1.22 (3H, t, J=7.0Hz), 3.06 (2H, d, J=6.2Hz), 3.11 (3H, s), 3.60-4.30 (11H, m), 6.40-6.80 (4H, m), 6.97 (1H, d, J=8.4Hz), 7.20-7.50 (6H, m), 8.00-8.20 (1H, m).

(4) The compound (488 mg) obtained in (3) above was dissolved in 5.0 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous sodium hydroxide solution (2.2 ml), and the mixture was stirred at room temperature for 6 hr. The solvent was evaporated under reduced pressure. Water (10 ml) and then saline were added to supersaturation, and the mixture was extracted three times with ethyl acetate (30 ml). The ethyl acetate layer was washed with saturated brine (10 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. Diethyl ether was added to the obtained residue. The solid was collected by filtration to give the title compound (356 mg).

IR ν (nujol) cm⁻¹; 1597, 1497.

¹H-NMR (MeOH-d₄) δ (ppm);

2.95-3.20 (2H, m), 3.07 (3H, s), 3.40-4.20 (9H, m), 6.40-6.70 (4H, m), 6.92 (1H, d, J=8.4Hz), 7.20-7.50 (3H, m), 7.90-8.15 (1H, m).

Example 12

Sodium 2-benzyl-7-[2-(5-ethyl-pyridin-2-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0096]

(1) Ethyl 2-benzyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.0 g) was dissolved in N,N-dimethylformamide (10 ml). Thereto was added sodium hydride (200 mg, 60% suspension in oil) under ice-cooling and the mixture was stirred at room temperature for 30 min to give solution (A). Separately, 5-ethyl-2-pyridineethanol (1.5 g) and triethylamine (1.68 ml) were dissolved in methylene chloride (40 ml). Thereto was added trifluoromethanesulfonic anhydride (2.0 ml) under ice-cooling, and the mixture was stirred at room temperature for 30 min. The reaction mixture was washed with water (30 ml) and dried over Na₂SO₄. Methylene chloride was evaporated under reduced pressure. The obtained 5-ethyl-2-pyridineethyl trifluoromethanesulfonate (2.81 g) was added to the above-mentioned solution (A), and the mixture was stirred at room temperature for 30 min. Ethyl acetate (100 ml) was added to the reaction mixture, and the mixture was washed with water (50 ml) and saturated brine (50 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-benzyl-7-[2-(5-ethyl-pyridin-2-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (0.58 g).

Ethyl 2-benzyl-7-[2-(5-ethyl-pyridin-2-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (nujol) cm⁻¹; 1732, 1612, 1504.

¹H-NMR (CDCl₃) δ (ppm);

1.22 (6H, t, J=7.2Hz), 2.61 (2H, q, J=7.2Hz), 3.07 (2H, d, J=5.5Hz), 3.18 (2H, t, J=6.6Hz), 3.72 (1H, t, J=5.5Hz),

3.81 (1H, s), 3.90 (4H, s), 4.13 (2H, q, J=7.2Hz), 4.27 (2H, t, J=6.6Hz), 6.51 (1H, d, J=2.0Hz), 6.69 (1H, dd, J=2.0, 8.4Hz), 6.98 (1H, d, J=8.4Hz), 7.10-7.50 (7H, m), 8.00-8.20 (1H, m).

(2) The compound (0.94 g) obtained in (1) above was dissolved in 40 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 2N aqueous sodium hydroxide solution (6.0 ml), and the mixture was stirred at 40°C for 2 hr. The solvent was evaporated under reduced pressure. Water (10 ml) and then sodium chloride were added to supersaturation, and the mixture was extracted three times with ethyl acetate (30 ml). The ethyl acetate layer was washed with saturated brine (10 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. Diisopropyl ether was added to the obtained residue. The solid was collected by filtration to give the title compound (0.58 g).

IR ν (nujol) cm⁻¹ ; 1576, 1504.

¹H-NMR (MeOH-d₄) δ (ppm);

1.22 (6H, t, J=7.5Hz), 2.63 (2H, q, J=7.5Hz), 2.90-3.20 (4H, m), 3.72 (1H, s), 3.85 (1H, s), 3.95-4.35 (5H, m), 4.27 (2H, t, J=6.6Hz), 6.40-6.75 (2H, m), 6.91 (1H, d, J=8.4Hz), 7.20-7.65 (7H, m), 8.20-8.35 (1H, m).

Example 13

2-Benzyl-7-[2-(indolin-1-yl)ethoxy]-1,2,3,4-tetrahydro-isoquinoline-(3S)-carboxylic acid

[0097]

(1) Ethyl 2-benzyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.0 g) was dissolved in N,N-dimethylformamide (10 ml). Thereto was added sodium hydride (154 mg, 60% suspension in oil) under ice-cooling and the mixture was stirred at the same temperature for 30 min. 1-(2-Bromoethyl)indoline (1.09 g) was added and the mixture was further stirred at room temperature for 2 hr. Ethyl acetate (100 ml) was added to the reaction mixture, and the mixture was washed with water (100 ml) and saturated brine (50 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-benzyl-7-[2-(indolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.18 g).

Ethyl 2-benzyl-7-[2-(indolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (neat) cm⁻¹ ; 2926, 2843, 1732, 1609, 1493.

¹H-NMR (CDCl₃) δ (ppm) ;

1.22 (3H, t, J=7.2Hz), 2.95 (2H, t, J=8.3Hz), 3.08 (2H, d, J=5.3Hz), 3.20-3.60 (3H, m), 3.60-4.30 (10H, m), 6.40-6.80 (4H, m), 6.80-7.20 (3H, m), 7.20-7.50 (5H, m).

(2) The compound (1.17 g) obtained in (1) above was dissolved in 24 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (7.69 ml), and the mixture was stirred at 50°C for 1 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (0.93 g).

IR ν (nujol) cm⁻¹ ; 1634, 1609, 1491.

¹H-NMR (CDCl₃) δ (ppm) ;

2.95 (2H, br-t, J=8.2Hz), 3.16 (2H, br-d, J=6.1Hz), 3.25-3.60 (3H, m), 3.60-4.40 (8H, m), 4.07 (2H, s), 4.10 (2H, t, J=5.5Hz), 4.13 (2H, q, J=7.2Hz), 5.60-6.30 (1H, br), 6.30-6.85 (4H, m), 6.85-7.20 (3H, m), 7.32 (5H, s).

Example 14

Ethyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0098] Ethyl 2-benzyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (10.0 g) and 2-(5-methyl-2-phenyloxazol-4-yl)ethyl methanesulfonate (18.01 g) were dissolved in N,N-dimethylformamide (200 ml). Thereto was added potassium carbonate (13.3 g) and the mixture was stirred at 80°C for 10 hr. Water (1 L) was added to the reaction mixture, and the mixture was extracted twice with ethyl acetate (200 ml). The ethyl acetate layer was washed with saturated brine (500 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give the title compound (7.06 g).

[0099] IR and NMR spectrum corresponded to those obtained in Example 2(1).

Example 15Methyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0100]

(1) Methyl 2-tert-butoxycarbonyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (15.16 g) and 2-(5-methyl-2-phenyloxazol-4-yl)ethyl methanesulfonate (20.0 g) were dissolved in N,N-dimethylformamide (300 ml). Thereto was added potassium carbonate (19.7 g) and the mixture was stirred at 80°C for 3.5 hr. Water (1 L) was added to the reaction mixture, and the mixture was extracted twice with ethyl acetate (300 ml). The ethyl acetate layer was washed with saturated brine (500 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give methyl 2-tert-butoxycarbonyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (14.0 g).

Methyl 2-tert-butoxycarbonyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

¹H-NMR (CDCl₃) δ (ppm);

1.46, 1.50 (9H, s, s), 2.36 (3H, s), 2.95 (2H, t, J=6.8Hz), 2.90-3.30 (2H, m), 3.60 (3H, s), 4.21 (2H, t, J=6.8Hz), 4.50, 4.60 (2H, s, s), 4.70-4.90, 5.00-5.20 (1H, m, m), 6.60-6.90 (2H, m), 7.12 (1H, d, J=8.4Hz), 7.30-7.55 (3H, m), 7.90-8.15 (2H, m).

(2) The compound (14.0 g) obtained in (1) above was dissolved in formic acid (42 ml). Thereto was added 8.78N hydrogen chloride solution (10.7 ml) in 2-propanol under ice-cooling, and the mixture was stirred at room temperature for 20 min. Ethyl acetate (300 ml) and water (500 ml) were added to the reaction mixture, and the mixture was neutralized with sodium hydrogencarbonate and the two layers were separated. The obtained ethyl acetate layer was washed with saturated brine (500 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give the title compound (9.4 g).

IR ν (nujol) cm⁻¹; 3560, 1744, 1643, 1612, 1578, 1553, 1504. ¹H-NMR (CDCl₃) δ (ppm);

1.92 (1H, s), 2.36 (3H, s), 2.80-3.20 (4H, m), 3.60-3.85 (1H, m), 3.76 (3H, s), 4.04 (2H, s), 4.21 (2H, t, J=6.6Hz), 6.57 (1H, d, J=2.0Hz), 6.71 (1H, dd, J=2.0, 8.6Hz), 7.00 (1H, d, J=8.6Hz), 7.30-7.60 (3H, m), 7.85-8.15 (2H, m).

Example 162-(4-Methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0101]

(1) The compound (800 mg) of Example 15 was dissolved in N,N-dimethylformamide (8 ml). Thereto was added sodium hydride (96 mg, 60% suspension in oil) under ice-cooling and the mixture was stirred at room temperature for 30 min. To the obtained mixture was dropwise added 4-methoxybenzyl chloride (0.41 ml), and the mixture was stirred further at 50°C for 3 hr. Ethyl acetate (50 ml) was added to the reaction mixture, and the mixture was washed with water (50 ml) and saturated brine (30 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give methyl 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (0.82 g). Methyl 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (nujol) cm⁻¹; 1736, 1638, 1614, 1553, 1514

¹H-NMR (CDCl₃) δ (ppm);

2.35 (3H, s), 2.93 (2H, t, J=6.6Hz), 3.05 (2H, d, J=5.5Hz), 3.66 (3H, s), 3.70-4.00 (8H, m), 4.17 (2H, t, J=6.6Hz), 6.50 (1H, d, J=2.0Hz), 6.68 (1H, dd, J=2.0, 8.6Hz), 6.85 (2H, d, J=8.6Hz), 6.98 (1H, d, J=8.6Hz), 7.28 (2H, d, J=8.6Hz), 7.30-7.50 (3H, m), 7.80-8.10 (2H, m).

(2) The compound (450 mg) obtained in (1) above was dissolved in 10 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 2N aqueous lithium hydroxide solution (2.5 ml), and the mixture was stirred at

room temperature for 3 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (350 mg).

IR ν (nujol) cm^{-1} ; 3288, 1612, 1555, 1514.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

2.35 (3H, s), 2.93 (2H, t, $J=6.4\text{Hz}$), 3.18 (2H, d, $J=6.8\text{Hz}$) 3.70-4.10 (5H, m), 3.77 (3H, s), 4.17 (2H, t, $J=6.4\text{Hz}$), 4.50 (1H, br-s), 6.60 (1H, d, $J=2.0\text{Hz}$), 6.65-6.95 (3H, m), 7.08 (2H, d, $J=8.4\text{Hz}$), 7.20-7.60 (5H, m), 7.80-8.10 (2H, m).

Example 17

Ethyl 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0102] In the same manner as in Example 16(1), the title compound was obtained.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.21 (3H, t, $J=7.0\text{Hz}$), 2.34 (3H, s), 2.91 (2H, t, $J=7.0\text{Hz}$), 3.04 (2H, d, $J=5.5\text{Hz}$), 3.60-3.95 (8H, m), 4.12 (2H, q, $J=7.0\text{Hz}$), 4.15 (2H, t, $J=7.0\text{Hz}$), 6.51 (1H, d, $J=2.0\text{Hz}$), 6.67 (1H, dd, $J=2.0, 8.8\text{Hz}$), 6.75-7.00 (3H, m), 7.15-7.50 (5H, m), 7.85-8.10 (2H, m).

Example 18

2-(4-Methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0103]

(1) The compound (800 mg) of Example 15 was dissolved in N,N-dimethylformamide (8.0 ml). Thereto was added sodium hydride (96 mg, 60% suspension in oil) under ice-cooling and the mixture was stirred at room temperature for 30 min. To the obtained mixture was dropwise added α -chloro-p-xylene (0.40 ml), and the mixture was stirred at 50°C for 3 hr and further stirred at room temperature for 15 hr. Ethyl acetate (50 ml) was added to the reaction mixture, and the mixture was washed with water (50 ml) and saturated brine (30 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give methyl 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (0.90 g).

Methyl 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (nujol) cm^{-1} ; 1736, 1639, 1614, 1595, 1551.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

2.34 (6H, s), 2.93 (2H, t, $J=6.7\text{Hz}$), 3.06 (2H, d, $J=5.0\text{Hz}$), 3.50-4.00 (5H, m), 3.65 (3H, s), 4.17 (2H, t, $J=6.7\text{Hz}$), 6.51 (1H, d, $J=2.0\text{Hz}$), 6.68 (1H, dd, $J=2.0, 8.6\text{Hz}$), 6.98 (2H, d, $J=8.6\text{Hz}$), 7.11 (2H, d, $J=8.4\text{Hz}$), 7.26 (2H, d, $J=8.4\text{Hz}$), 7.30-7.55 (3H, m), 7.80-8.15 (2H, m).

(2) The compound (608 mg) obtained in (1) above was dissolved in 17 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (6.1 ml), and the mixture was stirred at room temperature for 3 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (400 mg).

IR ν (nujol) cm^{-1} ; 1620, 1555, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

2.32 (3H, s), 2.35 (3H, s), 2.93 (2H, t, $J=7.0\text{Hz}$), 3.17 (2H, d, $J=6.6\text{Hz}$), 3.65-4.05 (5H, m), 4.17 (2H, t, $J=7.0\text{Hz}$), 4.73 (1H, br-s), 6.60 (1H, d, $J=2.0\text{Hz}$), 6.77 (1H, dd, $J=2.0, 8.8\text{Hz}$), 6.95-7.60 (8H, m), 7.85-8.10 (2H, m).

Example 19

Ethyl 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0104] In the same manner as in Example 18(1), the title compound was obtained.

¹H-NMR (CDCl₃) δ (ppm);

1.21 (3H, t, J=7.0Hz), 2.34 (3H, s), 2.92 (2H, t, J=7.0Hz), 3.05 (2H, d, J=5.4Hz), 3.71 (1H, t, J=5.4Hz), 3.80 (1H, s), 3.92 (1H, s), 4.12 (2H, q, J=7.0Hz), 4.16 (2H, t, J=7.0Hz), 6.51 (1H, d, J=2.0Hz), 6.68 (1H, dd, J=2.0, 8.4Hz), 6.98 (1H, d, J=8.4Hz), 7.00-7.60 (7H, m), 7.80-8.10 (2H, m).

Example 20

2-Benzyl-7-[2-(6-carboxyindolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0105]

(1) Ethyl 2-benzyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.0 g) was dissolved in N,N-dimethylformamide (10 ml). Thereto was added sodium hydride (154 mg, 60% suspension in oil) under ice-cooling and the mixture was stirred at the same temperature for 30 min. To the obtained mixture was added 1-(2-bromoethyl)-6-methoxycarbonylindoline (1.36 g), and the mixture was stirred further at room temperature for 4 hr. Ethyl acetate (100 ml) was added to the reaction mixture, and the mixture was washed with water (100 ml) and saturated brine (50 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give ethyl 2-benzyl-7-[2-(6-methoxycarbonylindolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.18 g).

Ethyl 2-benzyl-7-[2-(6-methoxycarbonylindolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (neat) cm⁻¹; 2949, 2841, 1717, 1611, 1587, 1497.

¹H-NMR (CDCl₃) δ (ppm);

1.23 (3H, t, J=7.2Hz), 2.99 (2H, t, J=8.8Hz), 3.00-3.25 (2H, m), 3.35-4.35 (13H, m), 3.87 (3H, s), 6.52 (1H, d, J=2.4Hz), 6.70 (1H, dd, J=2.4, 8.3Hz), 6.90-7.20 (3H, m), 7.20-7.55 (6H, m).

(2) The compound (1.31 g) obtained in (1) above was dissolved in 33 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (15.3 ml), and the mixture was stirred at 50°C for 2 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (1.0 g).

IR ν (nujol) cm⁻¹; 3400, 1693, 1612, 1497.

¹H-NMR (MeOH-d₄) δ (ppm);

2.96 (2H, br-t, J=8.0Hz), 3.15-3.75 (3H, m), 3.80-4.50 (10H, m), 6.65-7.70 (11H, m).

Example 21

2-(4-Fluorobenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0106]

(1) The compound (1.00 g) obtained in Example 15 was dissolved in N,N-dimethylformamide (20 ml). Thereto were added 4-fluorobenzyl chloride (0.46 ml), potassium carbonate (0.53 g) and potassium iodide (0.21 g), and the mixture was stirred at 50°C for 1.5 hr. Ethyl acetate (50 ml) was added to the reaction mixture, and the mixture was washed with water (100 ml) and saturated brine (100 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give methyl 2-(4-fluorobenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (0.92 g).

Methyl 2-(4-fluorobenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (nujol) cm⁻¹; 1738, 1639, 1616, 1551, 1510

¹H-NMR (CDCl₃) δ (ppm);

2.35 (3H, s), 2.93 (2H, t, J=6.8Hz), 3.07 (2H, d, J=5.0Hz), 3.55-4.00 (5H, m), 3.65 (3H, s), 4.18 (2H, t, J=6.8Hz), 6.51 (1H, d, J=2.0Hz), 6.70 (1H, dd, J=2.0, 8.6Hz), 6.80-7.15 (3H, m), 7.15-7.50 (5H, m), 7.80-8.15 (2H, m).

(2) The compound (900 mg) of (1) above was dissolved in 18 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (8.9 ml), and the mixture was stirred at room temperature for 3 hr. The solvent was evaporated under reduced pressure, and the obtained residue was acidified with citric acid. The precipitated crystals were collected by filtration to give the title compound (0.68 g).

IR ν (nujol) cm⁻¹; 3398, 1614, 1555, 1510.

¹H-NMR (CDCl₃) δ (ppm);

2.36 (3H, s), 2.94 (2H, t, J=6.4Hz), 3.15 (2H, d, J=6.4Hz), 3.45-4.00 (5H, m), 4.19 (2H, t, J=6.4Hz), 6.60 (1H, d, J=2.0Hz), 6.75 (1H, dd, J=2.0, 8.6Hz), 6.90-7.55 (8H, m), 7.90-8.10 (2H, m).

Example 22

2-(2,2-Dimethylpropionyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0107] Methyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (0.60 g) was dissolved in methylene chloride (6 ml). Thereto were added pivaloyl chloride (0.22 ml) and triethylamine (0.32 ml) under ice-cooling, and the mixture was stirred at the same temperature for 15 min. Ethyl acetate (100 ml) was added to the reaction mixture, and the mixture was washed successively with 10% aqueous citric acid solution (50 ml), saturated aqueous sodium hydrogencarbonate solution (50 ml) and saturated brine (50 ml) and dried over Na₂SO₄. The solvent was evaporated under reduced pressure. The obtained residue was dissolved in 18 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (4.6 ml), and the mixture was stirred at 50°C for 30 min. The solvent was evaporated under reduced pressure. Thereto was added water (20 ml) and the mixture was acidified with citric acid. The obtained mixture was extracted with ethyl acetate (50 ml). The ethyl acetate layer was washed with saturated brine (30 ml) was dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure to give the title compound (0.65 g).

IR ν (nujol) cm⁻¹; 1734, 1630, 1612, 1553.

¹H-NMR (DMSO-d₆) δ (ppm);

1.24 (9H, s), 2.36 (3H, s), 2.80-4.00 (1H, br), 2.92 (2H, t, J=6.4Hz), 2.95-3.15 (2H, m), 4.18 (2H, t, J=6.4Hz), 4.41, 4.91 (2H, ABq, J=18.1Hz), 4.90-5.15 (1H, m), 6.77 (1H, dd, J=2.0, 8.1Hz), 6.89 (1H, d, J=2.0Hz), 7.09 (1H, d, J=8.1Hz), 7.30-7.65 (3H, m), 7.80-8.10 (2H, m).

Example 23

2-(2,2-Dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0108] The title compound (1.66 g) of Example 22 was dissolved in pyridine (16.6 ml). Thereto was added sodium borohydride (1.36 g) and the mixture was stirred at 100°C for 4 hr. The mixture was acidified with 10% aqueous citric acid solution and extracted with ethyl acetate (100 ml). The ethyl acetate layer was washed successively by adding 10% aqueous citric acid solution (100 ml) and saturated brine (50 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give the title compound (0.84 g).

IR ν (nujol) cm⁻¹; 3391, 3279, 1668, 1645, 1616, 1597, 1497. ¹H-NMR (CDCl₃) δ (ppm);

0.96 (9H, s), 2.35 (3H, s), 2.46, 2.73 (2H, ABq, J=13.9Hz), 2.93 (2H, t, J=6.7Hz), 3.03-3.23 (2H, m), 3.57-3.78 (1H, m), 3.91, 4.18 (1H, ABq, J=15.4Hz), 4.17 (2H, t, J=6.7Hz), 5.60-6.05 (1H, br), 6.60 (1H, d, J=2.0Hz), 6.73 (1H, dd, J=2.0, 8.4Hz), 7.04 (1H, d, J=8.4Hz), 7.30-7.55 (3H, m), 7.80-8.10 (2H, m).

Example 24

2-(2,2-Dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

(1) Methyl 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0109] Methyl 2-(2,2-dimethylpropyl)-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.5 g) and 2-(5-methyl-2-phenyloxazol-4-yl)ethyl methanesulfonate (2.59 g) were dissolved in toluene (45 ml). Thereto were added potassium carbonate (2.24 g) and tetraethylammomium fluoride (0.60 g) and the mixture was stirred at 80°C for 3 hr. The reaction mixture was washed successively with water (50 ml) and saturated brine (50 ml) and dried over Na₂SO₄. Toluene was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to give methyl 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (2.77 g).

¹H-NMR (CDCl₃) δ (ppm) ;

0.88 (9H, s), 2.35, 2.58 (2H, ABq, J=14.5Hz), 2.36 (3H, s), 2.46 (2H, t, J=6.8Hz), 2.95-3.20 (2H, m), 3.60 (3H, s), 3.60-3.80 (1H, m), 3.85-4.20 (2H, m), 4.19 (2H, t, J=6.8Hz), 6.54 (1H, d, J=2.0Hz), 6.68 (1H, dd, J=2.0, 8.4Hz), 6.98 (1H, d, J=8.4Hz), 7.30-7.50 (3H, m), 7.85-8.05 (2H, m). (2) The compound (5.0 g) of (1) above was dissolved in 130 ml of a mixed solution of tetrahydrofuran - methanol (3:1). Thereto was added 1N aqueous lithium hydroxide solution (54 ml), and the mixture was stirred at 50°C for 3.5 hr and acidified with 6N hydrochloric acid. The solvent was evaporated under reduced pressure. The obtained residue was extracted with ethyl acetate (200 ml). The ethyl acetate layer was washed with saturated brine (100 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was dissolved in ethanol (25 ml). Thereto was added water (150 ml) and the mixture was stirred at room temperature to allow crystallization. The precipitated crystals were collected by filtration to give the title compound (4.52 g).

[0110] IR and ¹H-NMR NMR spectrum are correspondent to that obtained in Example 23.

Example 25

2-Benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid hydrochloride

[0111] The title compound (675 mg) of Example 2 was dissolved in 75% ethanol (10.1 ml) under heating. Thereto was added 6N hydrochloric acid (2.23 ml), and the mixture was stood at room temperature for 2 hr to allow crystallization. The precipitated crystals were collected by filtration to give the title compound (625 mg).

IR ν (nujol) cm⁻¹ ; 3398, 1734, 1680, 1641, 1620, 1587, 1574, 1551.

¹H-NMR (DMSO-d₆) δ (ppm) ;

2.36 (3H, s), 2.92 (2H, t, J=6.0Hz), 3.00-3.70 (2H, m), 4.19 (2H, t, J=6.0Hz), 4.25-4.75 (5H, m), 4.80-6.70 (2H, br), 6.70-7.05 (2H, m), 7.20 (1H, d, J=8.6Hz), 7.30-7.77 (8H, m), 7.80-8.10 (2H, m).

Example 26

Sodium 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0112] The title compound (1.0 g) of Example 2 was suspended in methanol (10 ml). Thereto was added a 2.09N sodium hydroxide solution (1.02 ml) in methanol. After dissolution, methanol was evaporated under reduced pressure. Diethyl ether was added to the obtained residue, and the precipitated crystals were collected by filtration to give the title compound (1.03 g).

IR ν (nujol) cm⁻¹ ; 1638, 1589, 1503.

¹H-NMR (DMSO-d₆) δ (ppm);

2.33 (3H, s), 2.55-3.60 (6H, m), 3.60-4.30 (5H, m), 6.47 (1H, s), 6.60 (1H, d, J=8.6Hz), 6.92 (1H, d, J=8.6Hz), 7.00-7.67 (8H, m), 7.67-8.05 (2H, m).

Example 27

2-(2,2-Dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid hydrochloride

[0113] The title compound (2.1 g) of Example 23 was dissolved in methanol (10.5 ml). 8.78N Hydrogen chloride solution (1.07 ml) in isopropanol and then ethyl acetate (50 ml) were added to the mixture and the mixture was stirred at room temperature for crystallization. The precipitated crystals were collected by filtration to give the title compound (1.03 g).

IR ν (nujol) cm^{-1} ; 3362, 3206, 1740, 1672, 1612, 1576, 1553. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;
 1.12 (9H, s), 2.37 (3H, s), 2.65-3.50 (6H, m),
 4.21 (2H, t, $J=6.5\text{Hz}$), 4.40-4.80 (3H, m), 4.85-6.50 (2H, br), 6.90 (1H, d, $J=8.1\text{Hz}$), 6.94 (1H, s), 7.21 (1H, d, $J=8.1\text{Hz}$), 7.35-7.65 (3H, m), 7.80-8.05 (2H, m).

Example 28

2-(2,2-Dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl) ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid sulfate

[0114] The title compound (0.5 g) of Example 23 was dissolved in methanol (1.25 ml). Sulfuric acid (0.3 ml) and then water (16.8 ml) were added to the mixture and the mixture was stirred at room temperature for crystallization. The precipitated crystals were collected by filtration to give the title compound (0.25 g).

IR ν (nujol) cm^{-1} ; 3400, 1715, 1650, 1615, 1550.
 $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;
 0.93 (9H, s), 2.36 (3H, s), 2.40-3.30 (6H, m), 3.75-4.45 (5H, m), 4.60-6.50 (2H, br), 6.70 (1H, br-s), 6.74 (1H, d, $J=8.1\text{Hz}$), 7.06 (1H, d, $J=8.1\text{Hz}$), 7.25-7.65 (3H, m), 7.75-8.10 (2H, m).

Example 29

2-(2,2-Dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid tosylate

[0115] The title compound (0.5 g) of Example 23 and p-toluenesulfonic acid (0.28 g) were dissolved in ethanol (10 ml) under heating. After crystallization at room temperature under stirring, the precipitated crystals were collected by filtration to give the title compound (0.3 g).

IR ν (nujol) cm^{-1} ; 3047, 1734, 1645, 1612, 1514,
 $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;
 1.07 (9H, s), 2.28 (3H, s), 2.36 (3H, s), 2.70-3.50 (6H, m), 4.21 (2H, t, $J=6.5\text{Hz}$), 4.40-4.80 (3H, m), 6.80-7.35 (5H, m), 7.35-7.65 (5H, m), 7.75-8.05 (2H, m), 8.40-12.00 (2H, br).

Example 30

2-(2,2-Dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid fumarate

[0116] The title compound (1.0 g) of Example 23 and fumaric acid (0.23 g) were dissolved in methanol (5 ml). Water (5 ml) was added to the mixture and the mixture was stirred at room temperature for crystallization. The precipitated crystals were collected by filtration to give the title compound (0.94 g).

IR ν (nujol) cm^{-1} ; 3500, 3395, 1680, 1650, 1625, 1575, 1550. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);
 0.85 (9H, s), 2.00-6.30 (2H, br), 2.35 (3H, s), 2.32, 2.59 (2H, ABq, $J=14.9\text{Hz}$), 2.75-3.10 (2H, m), 2.85 (2H, t, $J=6.6\text{Hz}$), 3.45-4.30 (3H, m), 4.15 (2H, t, $J=6.6\text{Hz}$), 6.61 (1H, d, $J=2.2\text{Hz}$), 6.64 (1H, s), 6.66 (1H, dd, $J=2.2, 8.4\text{Hz}$), 7.00 (1H, d, $J=8.4\text{Hz}$), 7.35-7.65 (3H, m), 7.75-8.10 (2H, m).

Example 31

Calcium 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0117] The title compound (0.9 g) of Example 23 was dissolved in ethanol (9 ml). 0.04N Aqueous calcium hydroxide solution (54 ml) was added to the mixture and the mixture was stirred at room temperature for crystallization. The precipitated crystals were collected by filtration to give the title compound (0.79 g).

IR ν (nujol) cm^{-1} ; 3396, 1638, 1611, 1556, 1504.

$^1\text{H-NMR}$ (MeOH-d_4) δ (ppm);

0.87 (9H, s), 2.34 (3H, s), 2.36, 2.58 (2H, ABq, $J=14.0\text{Hz}$), 2.80-3.10 (2H, m), 2.90 (2H, t, $J=6.5\text{Hz}$), 3.30-3.80 (3H, m), 4.15 (2H, t, $J=6.5\text{Hz}$), 6.50 (1H, d, $J=2.4\text{Hz}$), 6.60 (1H, dd, $J=2.4, 8.2\text{Hz}$), 6.90 (1H, d, $J=8.2\text{Hz}$), 7.30-7.60 (3H, m), 7.80-8.05 (2H, m).

[0118] The following compounds were synthesized in the same manner as in Examples 1-24.

Example 32

2-Benzyl-7-[2-(5-methyl-2-tert-butyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0119]

IR ν (nujol) cm^{-1} ; 3458, 1682, 1618, 1587, 1510.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

1.32 (9H, s), 2.23 (3H, s), 2.83 (2H, t, $J=6.6\text{Hz}$), 3.18 (2H, d, $J=5.9\text{Hz}$), 3.65-4.40 (7H, m), 5.60 (1H, br-s), 6.56 (1H, br-s), 6.73 (1H, br-d), 7.06 (1H, d, $J=8.4\text{Hz}$), 7.20-7.55 (5H, m).

Example 33

2-Benzyl-7-[2-(5-methyl-2-(thiophen-2-yl)oxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0120]

IR ν (nujol) cm^{-1} ; 3423, 1616, 1578, 1510.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

2.31 (3H, s), 2.70-3.10 (4H, m), 3.40-4.00 (4H, m), 3.39 (2H, s), 4.11 (2H, d, $J=6.2\text{Hz}$), 6.59 (1H, br-s), 6.67 (1H, d, $J=8.4\text{Hz}$), 7.01 (1H, d, $J=8.4\text{Hz}$), 7.05-7.80 (3H, m), 7.32 (5H, s).

Example 34

2-Benzyl-7-[2-(5-methyl-2-isopropylloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0121]

IR ν (nujol) cm^{-1} ; 3456, 1684, 1614, 1576, 1510.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);

1.21 (6H, m), 2.19 (3H, s), 2.70-3.10 (5H, m), 3.50-4.20 (5H, m), 6.40-6.85 (2H, m), 7.01 (1H, d, $J=8.1\text{Hz}$), 7.34 (5H, br-s).

Example 35

2-Butyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0122]

IR ν (nujol) cm^{-1} ; 3382, 1722, 1614, 1554, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

0.88 (3H, t, J=6.6Hz), 1.10-1.95 (4H, m), 2.36 (3H, s), 2.75-3.40 (6H, m), 3.71 (2H, br-t), 3.95-4.25 (4H, m), 6.57-7.57 (6H, m), 7.80-8.10 (3H, m).

Example 36

2-Benzyl-7-(2-[5-methyl-2-(2-methylpropenyl)oxazol-4-yl]ethoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0123]

IR v (nujol) cm^{-1} ; 3443, 3300, 1695, 1655, 1622, 1543, 1508. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;
1.89 (3H, s), 2.11 (3H, s), 2.27 (3H, s), 2.79 (2H, t, J=6.1Hz), 2.90-3.20 (2H, m), 3.50-4.00 (4H, m), 3.93 (2H, s),
4.07 (2H, t, J=6.1Hz), 5.99 (1H, s), 6.58 (1H, s), 6.67 (1H, d, J=8.2Hz), 6.72 (1H, d, J=8.2Hz), 7.33 (5H, s).

Example 37

2-Benzyl-7-(2-[2-(3-butenyl)-5-methyloxazol-4-yl]ethoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0124]

IR v (nujol) cm^{-1} ; 3442, 1688, 1614, 1578, 1508.
 $^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;
2.23 (3H, s), 2.49 (2H, t, J=6.2Hz), 2.65-2.90 (4H, m), 3.05-3.30 (2H, m), 3.75-4.50 (8H, m), 4.90-5.20 (2H, m),
5.65-6.10 (1H, m), 6.58 (1H, d, J=1.7Hz), 6.75 (1H, dd, J=1.7, 8.2Hz), 7.07 (1H, d, J=8.2Hz), 7.35 (5H, s).

Example 38

2-Allyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0125]

IR v (nujol) cm^{-1} ; 3335, 1690, 1618, 1553, 1506.
 $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;
2.35 (3H, s), 2.70-3.15 (4H, m), 3.38 (2H, d, J=6.2Hz), 3.55-4.00 (3H, m), 4.16 (2H, t, J=6.6Hz), 4.40-5.50 (1H, br)
5.00-5.40 (2H, m), 5.60-6.10 (1H, m), 6.65 (1H, s), 6.69 (1H, d, J=8.1Hz), 7.01 (1H, d, J=8.1Hz), 7.35-7.65 (3H, m),
7.75-8.10 (2H, m).

Example 39

7-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-(2-propynyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0126]

IR v (nujol) cm^{-1} ; 3383, 3306, 3221, 1692, 1622, 1508.
 $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;
2.00-6.40 (1H, br), 2.35 (3H, s), 2.70-3.10 (4H, m), 3.10-3.25 (1H, m), 3.50-4.00 (5H, m), 4.17 (2H, t, J=6.4Hz),
6.66 (1H, s), 6.70 (1H, d, J=8.6Hz), 7.01 (1H, d, J=8.6Hz), 7.30-7.70 (3H, m), 7.85-8.05 (2H, m).

Example 40

2-(2-Butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0127]

IR v (nujol) cm^{-1} ; 3447, 3342, 1684, 1620, 1556.
 $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;
1.67 (3H, d, J=4.9Hz), 2.35 (3H, s), 2.70-3.10 (4H, m), 3.20-3.50 (2H, m), 3.50-4.00 (3H, m), 4.16 (2H, t, J=6.4Hz),
4.35-5.20 (1H, br), 5.25-5.90 (2H, m), 6.55-6.90 (2H, m), 7.01 (2H, d, J=8.1Hz), 7.35-7.70 (3H, m), 7.75-8.10 (2H, m).

Example 412-Benzyl-7-[(indolin-3-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0128]

IR v (nujol) cm^{-1} ; 1611, 1506. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

1.60-2.30 (2H, m), 2.80-4.20 (10H, m), 3.91 (2H, s), 4.20-6.00 (1H, br), 6.25-7.10 (7H, m), 7.33 (5H, s).

Example 422-(3-Butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0129]

IR v (nujol) cm^{-1} ; 3425, 1682, 1612, 1555. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;2.10-2.40 (2H, m), 2.35 (3H, s), 2.60-3.15 (6H, m), 3.50-4.00 (3H, m), 4.17 (2H, t, $J=6.3\text{Hz}$), 4.40-5.40 (1H, br), 4.85-5.25 (2H, m), 5.55-6.10 (1H, m), 6.50-6.85 (2H, m), 7.01 (1H, d, $J=8.1\text{Hz}$), 7.35-7.70 (3H, m), 7.75-8.05 (2H, m).**Example 43**7-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-pentanoyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0130]

IR v (nujol) cm^{-1} ; 1742, 1639, 1611, 1572, 1506. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;0.88 (3H, br-t), 1.05-1.75 (4H, m), 2.15-2.70 (2H, m), 2.35 (3H, s), 2.70-3.30 (4H, m), 4.18 (2H, br-t), 4.30-4.90 (2H, m), 4.90-5.25 (1H, m), 6.60-6.95 (2H, m), 7.08 (1H, d, $J=7.9\text{Hz}$), 7.35-7.70 (3H, m), 7.75-8.10 (2H, m), 11.00-13.00 (1H, br).**Example 44**7-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-(4-pentenyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0131]

IR v (nujol) cm^{-1} ; 1742, 1641, 1611, 1570. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;2.15-2.70 (2H, m), 2.35 (3H, s), 2.70-3.30 (4H, m), 4.18 (2H, br-t), 4.37-5.50 (5H, m), 5.60-6.15 (1H, m), 6.60-6.95 (2H, m), 7.09 (1H, d, $J=7.7\text{Hz}$), 7.30-7.75 (3H, m), 7.75-8.15 (2H, m), 11.00-13.00 (1H, br).**Example 45**2-(3-Methyl-2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0132]

IR v (nujol) cm^{-1} ; 1738, 1641, 1611, 1555. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;1.87 (6H, s), 2.35 (3H, s), 2.70-3.30 (4H, m), 4.18 (2H, br-t), 4.49 (1H, d, $J=18.0\text{Hz}$), 4.76 (1H, d, $J=18.0\text{Hz}$), 4.95-5.22 (1H, m), 5.75-6.10 (1H, m), 6.50-6.90 (2H, m), 7.08 (1H, d, $J=7.5\text{Hz}$), 7.20-7.60 (3H, m), 7.60-8.05 (2H, m), 11.00-13.00 (1H, br).

Example 46

2-(3,3-Dimethylbutynyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0133]

IR ν (nujol) cm^{-1} ; 1738, 1639, 1611, 1583, 1555.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

1.02 (9H, s), 2.36 (5H, s), 2.73-3.20 (4H, m), 4.17 (2H, t, $J=7.0\text{Hz}$), 4.50 (1H, d, $J=9.0\text{Hz}$), 4.83 (1H, d, $J=9.0\text{Hz}$), 5.12 (1H, t, $J=6.0\text{Hz}$), 6.60-6.95 (2H, m), 7.10 (1H, d, $J=7.0\text{Hz}$), 7.35-7.65 (3H, m), 7.80-8.05 (2H, m), 11.00-13.00 (1H, br).

Example 47

2-Benzyl-7-methoxy-6-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3RS)-carboxylic acid

[0134]

IR ν (nujol) cm^{-1} ; 1722, 1628, 1553, 1520.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);

2.36 (3H, s), 3.00 (2H, t, $J=6.8\text{Hz}$), 3.10-3.35 (2H, m), 3.80-4.10 (3H, m), 3.75 (3H, s), 4.23 (2H, t, $J=6.8\text{Hz}$), 5.80-6.20 (1H, br), 6.50, 6.72 (2H, s, s), 7.20-7.60 (8H, m), 7.80-8.10 (2H, m).

Example 48

Sodium 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(pyridin-2-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0135]

IR ν (nujol) cm^{-1} ; 1609, 1575, 1554, 1502.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

2.32 (3H, s), 2.60-3.20 (4H, m), 3.20-3.90 (5H, m), 4.08 (2H, br-t, $J=6.5\text{Hz}$), 6.15-6.40 (1H, m), 6.40-6.70 (1H, m), 7.75-8.20 (3H, m), 7.20-7.65 (4H, m), 7.75-8.10 (2H, m), 8.25-8.60 (1H, m).

Example 49

2-Benzyl-7-(3-methyl-3-phenylbutoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0136]

IR ν (nujol) cm^{-1} ; 1612, 1506.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

1.32 (6H, s), 2.02 (2H, t, $J=7.5\text{Hz}$), 2.80-3.10 (2H, m), 3.40-4.00 (5H, m), 3.88 (2H, s), 4.10-6.00 (1H, br), 6.36 (1H, d, $J=2.0\text{Hz}$), 6.53 (1H, dd, $J=2.0, 8.6\text{Hz}$), 6.96 (1H, d, $J=8.6\text{Hz}$), 7.10-7.55 (10H, m).

Example 50

2-Benzyl-7-(3,3-dimethyl-4-phenylbutoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0137]

IR ν (nujol) cm^{-1} ; 1611, 1506.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);

0.87 (6H, s), 1.61 (2H, t, $J=7.0\text{Hz}$), 2.54 (2H, s), 2.85-3.15 (2H, m), 3.50-4.20 (5H, m), 3.91 (2H, s), 4.20-6.00 (1H, br), 6.60 (1H, br-s), 6.67 (1H, d, $J=8.6\text{Hz}$), 7.01 (1H, d, $J=8.6\text{Hz}$), 7.05-7.50 (10H, m).

Example 512-Benzyl-7-(2-isopropylbenzoxazol-6-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid**[0138]**IR v (Nujol) cm^{-1} ; 1632, 1585, 1572, 1501. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);1.37 (6H, d, $J=7.0\text{Hz}$), 2.85-3.45 (4H, m), 3.50-4.20 (6H, m), 5.12 (2H, s), 6.69 (1H, s), 6.77 (1H, d, $J=8.6\text{Hz}$), 7.04 (1H, d, $J=8.6\text{Hz}$), 7.25-7.85 (8H, m).**Example 52**2-Benzyl-7-(2-tert-butylbenzoxazol-6-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid**[0139]**IR v (Nujol) cm^{-1} ; 1611, 1583, 1562, 1506. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;1.42 (9H, s), 2.85-3.10 (2H, br), 3.50-4.20 (6H, m), 5.13 (2H, s), 6.70 (1H, s), 6.77 (1H, d, $J=8.4\text{Hz}$), 7.04 (1H, d, $J=8.4\text{Hz}$), 7.20-7.50 (6H, m), 7.55-7.65 (1H, m), 7.70 (1H, s).**Example 53**2-Benzyl-7-(2-tert-butylbenzoxazol-5-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid**[0140]**IR v (nujol) cm^{-1} ; 1717, 1645, 1612, 1553. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);1.43 (9H, s), 2.85-3.15 (2H, m), 3.50-4.15 (3H, m), 3.90 (2H, s), 5.11 (2H, s), 6.69 (1H, br-s), 6.75 (1H, d, $J=8.1\text{Hz}$), 7.03 (1H, d, $J=8.1\text{Hz}$), 7.10-7.50 (6H, m), 7.65 (1H, d, $J=9.0$), 7.71 (1H, br-s).**Example 54**7-(2-tert-Butylbenzoxazol-6-yl)methoxy-2-(2,2-dimethylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)- carboxylic acid**[0141]**IR v (nujol) cm^{-1} ; 1740, 1612, 1560, 1508. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;1.10 (9H, s), 1.43 (9H, s), 2.79, 3.19 (2H, ABq, $J=13.6\text{Hz}$), 3.20-3.45 (3H, m), 4.25-4.55 (4H, br), 5.20 (2H, s), 6.96 (1H, d, $J=8.1\text{Hz}$), 7.00 (1H, s), 7.20 (1H, d, $J=8.1\text{Hz}$), 7.40 (1H, d, $J=8.1\text{Hz}$), 7.68 (1H, d, $J=8.1\text{Hz}$) 7.73 (1H, s).**Example 55**2-Benzyl-7-(2-isopropylbenzoxazol-5-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid**[0142]**IR v (nujol) cm^{-1} ; 1634, 1587, 1570, 1501. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);1.37 (6H, d, $J=6.8\text{Hz}$), 2.90-3.15 (2H, m), 3.25 (1H, quintet, $J=6.8\text{Hz}$), 3.50-4.30 (7H, m), 5.11 (2H, s), 6.69 (1H, s), 6.76 (1H, d, $J=8.1\text{Hz}$), 7.04 (1H, d, $J=8.1\text{Hz}$), 7.20-7.50 (6H, m), 7.55-7.65 (1H, m), 7.70 (1H, s).

Example 56

Sodium 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(pyridin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0143]

IR ν (nujol) cm^{-1} ; 3420, 3177, 1639, 1558, 1504.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);

2.34 (3H, s), 2.70-3.05 (4H, m), 3.10-3.60 (3H, m), 3.98 (2H, br-t, $J=5.7\text{Hz}$), 4.10-4.25 (2H, m), 6.51 (1H, br-s), 6.61 (1H, br-d, $J=8.7\text{Hz}$), 6.94 (1H, br-d, $J=8.7\text{Hz}$), 7.25-7.65 (5H, m), 7.75-8.00 (2H, m), 8.46 (2H, d, $J=5.2\text{Hz}$).

Example 57

Sodium 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-[(pyridin-2-yl)carbonyl]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0144]

IR ν (nujol) cm^{-1} ; 3385, 1624, 1566, 1504.

$^1\text{H-NMR}$ (MeOH-d_4) δ (ppm) ;

2.31, 2.36 (3H, s, s), 2.75-3.05 (2H, m), 3.05-3.30 (2H, m), 4.00-4.30 (2H, m), 4.50-5.30 (3H, m), 6.60-6.80 (2H, m) 7.03 (1H, dd, $J=2.0, 8.5\text{Hz}$), 7.30-7.75 (5H, m), 7.75-8.10 (3H, m), 8.50-8.70 (1H, m).

Example 58

Methyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0145]

IR ν (nujol) cm^{-1} ; 1736, 1639, 1612, 1504.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

2.34 (3H, s), 2.92 (2H, t, $J=7.0\text{Hz}$), 3.07 (2H, d, $J=5.0\text{Hz}$), 3.64 (3H, s), 3.64-4.00 (5H, m), 4.17 (2H, t, $J=7.0\text{Hz}$), 6.51 (1H, d, $J=2.0\text{Hz}$), 6.68 (1H, dd, $J=2.0, 8.4\text{Hz}$), 6.98 (1H, d, $J=8.4\text{Hz}$), 7.20-7.60 (8H, m), 7.80-8.10 (2H, m).

Example 59

2-Benzyl-7-[2-(2-cyclopropyl-5-methyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0146]

IR ν (nujol) cm^{-1} ; 3470, 1684, 1618, 1583, 1510.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

0.70-1.10 (4H, m), 1.80-2.20 (1H, m), 2.16 (3H, s), 2.60-2.85 (2H, m), 2.90-3.15 (2H, m), 3.50-4.20 (5H, m), 6.50-6.80 (2H, m), 7.03 (1H, d, $J=8.1\text{Hz}$), 7.34 (5H, s).

Example 60

2-(3-Methyl-2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0147]

IR ν (nujol) cm^{-1} ; 3447, 3335, 1670, 1668, 1622, 1556, 1506.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);

1.61 (3H, s), 1.72 (3H, s), 2.35 (3H, s), 2.70-3.20 (4H, m), 3.39 (2H, d, $J=7.0\text{Hz}$), 3.50-4.01 (3H, m), 4.16 (2H, t, $J=7.0\text{Hz}$), 4.35-5.60 (1H, br), 5.25 (1H, br-t), 6.67 (1H, s), 6.71 (1H, d, $J=8.4\text{Hz}$), 7.02 (1H, d, $J=8.4\text{Hz}$), 7.30-7.70 (8H, m), 7.75-8.10 (2H, m).

Example 61

2-(2,2-Dimethylpropyl)-7-[2-(5-methyl-2-tert-butyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0148]

IR v (nujol) cm^{-1} ; 1717, 1614, 1566, 1504.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

0.97 (9H, s), 1.33 (9H, s), 2.24 (3H, s), 2.44, 2.68 (2H, ABq, $J=13.9\text{Hz}$), 2.84 (2H, t, $J=6.7\text{Hz}$), 3.00-3.22 (2H, m), 3.65 (3H, t, $J=6.1\text{Hz}$), 3.83, 4.08 (2H, ABq, $J=15.1\text{Hz}$), 4.07 (2H, t, $J=6.7\text{Hz}$), 6.58 (1H, d, $J=1.7\text{Hz}$), 6.72 (1H, dd, $J=1.7, 8.4\text{Hz}$), 7.05 (1H, d, $J=8.4\text{Hz}$), 7.50-8.20 (1H, br).

Example 62

2-Benzyl-7-[2-(1-butenyl)-5-methyloxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0149]

IR v (nujol) cm^{-1} ; 3470, 1682, 1614, 1585, 1512.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

1.07 (3H, t, $J=7.5\text{Hz}$), 2.05-2.20 (1H, m), 2.26 (3H, s), 2.50-3.00 (3H, m), 3.65-4.45 (7H, m), 5.92 (1H, br-s), 6.17 (1H, d, $J=16.3\text{Hz}$), 6.45-6.85 (3H, m), 7.05 (1H, d, $J=8.4\text{Hz}$), 7.34 (5H, s).

Example 63

2-Benzyl-7-[2-(2,2-dimethylpropyl)-5-methyloxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0150]

IR v (nujol) cm^{-1} ; 1722, 1614, 1568, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

0.95 (9H, s), 2.34 (3H, s), 2.55 (2H, s), 2.60-3.00 (2H, m), 3.00-3.30 (2H, m), 3.80-4.40 (7H, m), 6.64 (1H, br-s), 6.70 (1H, d, $J=8.8\text{Hz}$), 7.02 (1H, d, $J=8.8\text{Hz}$), 7.32 (5H, s), 7.80 (1H, br-s).

Example 64

Ethyl 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate hydrochloride

[0151]

IR v (nujol) cm^{-1} ; 3400, 1744, 1676, 1614, 1589, 1574, 1553, 1508.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

1.12 (9H, s), 1.23 (3H, t, $J=7.0\text{Hz}$), 2.36 (3H, s), 2.60-3.60 (6H, m), 4.00-4.40 (4H, m), 4.40-6.00 (4H, m), 6.88 (1H, d, $J=8.0\text{Hz}$), 6.92 (1H, s), 7.18 (1H, d, $J=8.0\text{Hz}$), 7.35-7.70 (8H, m), 7.75-8.10 (2H, m).

Example 65

7-(Benzofran-2-ylmethoxy)-2-benzyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0152]

IR v (nujol) cm^{-1} ; 1632, 1587, 1501.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);

2.00-6.50 (1H, br), 2.85-3.15 (2H, m), 3.50-4.10 (3H, m), 3.91 (2H, s), 5.16 (2H, s), 6.60-7.80 (8H, m), 7.33 (5H, s).

Example 662-Isobutryl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0153]

IR v (nujol) cm^{-1} ; 1736, 1639, 1612, 1504. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;1.02 (6H, br-d), 2.35 (3H, s), 2.65-3.30 (5H, m), 4.00-5.30 (6H, m), 6.60-6.95 (2H, m), 7.09 (1H, d, $J=8.0\text{Hz}$),
7.25-7.70 (8H, m), 7.70-8.10 (2H, m).**Example 67**7-[2-(Benzofran-2-yl)ethoxy]-2-benzyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0154]

IR v (nujol) cm^{-1} ; 1634, 1585, 1501. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

2.80-3.10 (2H, m), 3.19 (2H, br-t), 3.45-4.10 (3H, m), 3.90 (2H, s), 4.25 (2H, br-t), 6.50-7.80 (9H, m), 7.33 (5H, s).

Example 687-[2-(5-Ethylpyridin-2-yl)ethoxy]-2-hexanoyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid hydrochloride

[0155]

 $^1\text{H-NMR}$ (CDCl_3) δ (ppm);0.87 (3H, br-t), 1.05-1.85 (9H, m), 2.15-2.55 (4H, m), 2.55-3.75 (4H, m), 4.00-4.90 (4H, m), 5.25-5.50 (1H, m),
6.40-7.10 (4H, m), 7.75 (1H, br-d), 8.15 (1H, br-d), 8.52 (1H, br-s).**Example 69**2-Carboxymethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0156]

IR v (nujol) cm^{-1} ; 1620, 1585, 1556, 1508. $^1\text{H-NMR}$ (DMSO-d_6) δ (ppm);2.35 (3H, s), 2.70-3.15 (4H, m), 3.41, 3.65 (2H, ABq, $J=17.5\text{Hz}$),
3.70-4.00 (3H, m), 4.16 (2H, t, $J=7.0\text{Hz}$), 6.00-11.00 (1H, br), 6.64 (1H, s), 6.69 (1H, d, $J=8.2\text{Hz}$), 7.01 (1H, d, $J=8.2\text{Hz}$), 7.20-7.70 (8H, m), 7.70-8.05 (2H, m).**Example 70**2-[3-(Methoxycarbonyl)propionyl]-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0157]

IR v (nujol) cm^{-1} ; 1732, 1652, 1554, 1505. $^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;2.33 (3H, s), 2.50-3.40 (7H, m), 3.65 (3H, s), 4.07 (2H, br-t), 4.45-5.50 (3H, m), 5.60-6.20 (1H, br), 6.59 (1H, br-s), 6.67 (1H, d, $J=8.0\text{Hz}$), 7.03 (1H, d, $J=8.0\text{Hz}$), 7.20-7.60 (3H, m), 7.80-8.10 (2H, m).

Example 71

2-[3-(Ethoxycarbonyl)propyl]-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0158]

IR v (nujol) cm^{-1} ; 3375, 1733, 1620, 1555, 1505.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.18 (3H, t, $J=7.0\text{Hz}$), 1.76-2.15 (2H, m), 2.15-2.50 (2H, m), 2.35 (3H, s), 2.70-3.35 (6H, m), 3.60-4.40 (7H, m), 5.27 (1H, br-s), 6.61 (1H, br-s), 6.73 (1H, d, $J=8.4\text{Hz}$), 7.03 (1H, d, $J=8.4\text{Hz}$), 7.25-7.55 (3H, m), 7.80-8.10 (2H, m).

Example 72

2-Benzyl-6-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3RS)-carboxylic acid

[0159]

IR v (nujol) cm^{-1} ; 1634, 1614, 1499.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

2.35 (3H, s), 2.65-3.25 (4H, m), 3.40-4.00 (3H, m), 3.90 (2H, s), 4.17 (2H, br-t), 6.20-10.00 (1H, br), 6.50-7.00 (2H, m), 6.71 (1H, s), 7.30-7.70 (3H, m), 7.32 (5H, s), 7.75-8.15 (2H, m).

Example 73

2-(3-Acetylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0160]

IR v (nujol) cm^{-1} ; 1682, 1620, 1508.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

2.33 (3H, s), 2.56 (3H, s), 2.70-3.20 (4H, m), 3.50-4.30 (5H, m), 3.97 (2H, s), 6.50-6.90 (2H, m), 7.02 (1H, d, $J=8.4\text{Hz}$), 7.30-8.00 (9H, m).

Example 74

2-(2-Acetylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0161]

IR v (nujol) cm^{-1} ; 1668, 1643, 1614, 1504.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

2.33 (3H, s), 2.36 (3H, s), 2.70-3.20 (4H, m), 3.30-4.30 (5H, m), 6.57 (1H, d, $J=2.0\text{Hz}$), 6.66 (1H, dd, $J=2.0, 8.4\text{Hz}$), 7.00 (1H, d, $J=8.4\text{Hz}$), 7.20-7.75 (7H, m), 7.75-8.10 (2H, m).

Example 75

2-Benzyl-7-[(5-methyl-2-phenyloxazol-4-yl)methoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid

[0162]

IR v (nujol) cm^{-1} ; 3462, 1680, 1614, 1556, 1508.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm) ;

2.41 (3H, s), 2.83-3.20 (2H, m), 3.44-4.20 (5H, m), 4.91 (2H, s), 6.73 (1H, br-s), 6.77 (1H, d, $J=8.1\text{Hz}$), 7.34 (1H, d, $J=8.1\text{Hz}$), 7.34 (5H, s), 7.40-7.68 (3H, m), 7.75-8.10 (2H, m).

Reference Example 1

Ethyl 2-tert-butoxycarbonyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0163]

(1) 3,5-Diiodo-L-tyrosine dihydrate (25.0 g) was suspended in concentrated hydrochloric acid (250 ml). Thereto were successively added 1,2-dimethoxyethane (18 ml) and 37% formalin (20 ml), and the mixture was warmed to 75°C over 30 min. To the reaction mixture were further added concentrated hydrochloric acid (120 ml), 1,2-dimethoxyethane (9 ml) and 37% formalin (10 ml), and the mixture was stirred at 75°C for 18 hr. The precipitated crystals were collected by filtration and washed with 1,2-dimethoxyethane (20 ml) to give 7-hydroxy-6,8-diiodo-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid hydrochloride (12.8 g).

7-Hydroxy-6,8-diiodo-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid hydrochloride:

IR ν (nujol) cm^{-1} ; 1751, 1599, 1578.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

3.00-3.30 (2H, m), 4.05 (2H, s), 4.30 (1H, dd, $J=5.9, 9.5\text{Hz}$), 7.71 (1H, s).

(2) The compound (12.8 g) obtained in (1) above was suspended in ethanol (500 ml). Thereto was added concentrated hydrochloric acid (10 ml) and the mixture was refluxed for 15 hr. Ethanol was evaporated under reduced pressure. Ethyl acetate (300 ml) was added to the obtained residue, and the compound was washed with saturated aqueous sodium hydrogencarbonate solution (100 ml) and saturated brine (100 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure to give ethyl 7-hydroxy-6,8-diiodo-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (11.11 g).

Ethyl 7-hydroxy-6,8-diiodo-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.29 (3H, t, $J=7.0\text{Hz}$), 2.80-3.00 (2H, m), 3.30-4.10 (5H, m), 4.23 (2H, q, $J=7.0\text{Hz}$), 7.46 (1H, s).

(3) 10% Pd-C (350 mg) was suspended in methanol (60 ml). Thereto were added triethylamine (2.0 ml) and the compound (2.80 g) obtained in (2) above, and the compound was catalytically hydrogenated at room temperature, 29.4×10^4 Pa (3.0 kgf/cm²) for 3 hr. Pd-C was filtered off, and methanol was evaporated under reduced pressure. Ethyl acetate (10 ml) was added to the obtained residue. The mixture was washed with saturated brine (100 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure to give ethyl 7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate (1.14 g).

Ethyl 7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate:

IR ν (nujol) cm^{-1} ; 1732, 1607, 1516.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

1.28 (3H, t, $J=7.0\text{Hz}$), 2.80-3.10 (3H, m), 3.60-3.80 (1H, m), 3.97 (2H, s), 4.05-4.20 (4H, m), 6.43 (1H, s), 6.50-6.80 (1H, m), 6.92 (1H, d, $J=8.4\text{Hz}$).

(4) The compound (1.13 g) obtained in (3) above was dissolved in tetrahydrofuran (20 ml). Thereto was added di-tert-butyl dicarbonate (1.50 g) and the mixture was stirred at room temperature for 1 hr. Ethyl acetate (30 ml) was added to the reaction mixture, and the mixture was washed with saturated brine (20 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by column chromatography to give the title compound (1.51 g).

IR ν (nujol) cm^{-1} ; 3260, 1756, 1671, 1615, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.29 (3H, t, $J=7.0\text{Hz}$), 1.47 (9H, s), 3.08 (2H, d, $J=5.2\text{Hz}$), 4.21 (2H, q, $J=7.0\text{Hz}$), 4.41 (1H, d, $J=15.5\text{Hz}$), 4.60-5.25 (1H, m), 4.65 (1H, d, $J=15.5\text{Hz}$), 5.00-6.00 (1H, br), 6.50-6.80 (2H, m), 6.98 (1H, d, $J=8.1\text{Hz}$).

Reference Example 2

Methyl 2-tert-butoxycarbonyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0164] In the same manner as in Reference Example 1, the title compound was obtained.

IR ν (nujol) cm^{-1} ; 3261, 1755, 1672, 1614, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.47 (9H, s), 3.08 (2H, d, $J=5.2\text{Hz}$), 3.63 (3H, s), 4.40 (1H, d, $J=16.5\text{Hz}$), 4.60-5.25 (1H, m), 4.66 (1H, d, $J=16.5\text{Hz}$), 5.60-6.60 (1H, br), 6.50-6.80 (2H, m), 6.99 (1H, d, $J=8.1\text{Hz}$).

Reference Example 3

2-(5-Methyl-2-phenyl-oxazol-4-yl)ethyl methanesulfonate

[0165] To methylene chloride (200 ml) were added 2-(5-methyl-2-phenyl-oxazol-4-yl)ethanol (20 g) and triethylamine (19.2 ml). Thereto was dropwise added methanesulfonyl chloride. (9.52 ml) at 0°C and the mixture was stirred at the same temperature for 15 min. The mixture was washed with 10% aqueous citric acid solution (200 ml), saturated aqueous sodium hydrogencarbonate solution (100 ml) and saturated brine (100 ml) and dried over Na_2SO_4 . Methylene chloride was evaporated under reduced pressure. The obtained residue was purified by column chromatography to give the title compound (21.45 g).

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

2.53 (3H, s), 2.94 (3H, s), 2.94 (2H, t, $J=7.0\text{Hz}$), 4.52 (2H, t, $J=7.0\text{Hz}$), 7.30-7.50 (3H, m), 7.80-8.10 (2H, m).

Reference Example 4

Ethyl 2-benzyl-7-hydroxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate

[0166] The compound (8.1 g) obtained in Reference Example 1(3) was dissolved in N,N -dimethylformamide (80 ml). Thereto were added triethylamine (2.0 ml) and benzyl bromide (4.57 ml) and the mixture was stirred at room temperature for 3 hr. Water (500 ml) was added to the reaction mixture, and the mixture was extracted twice with ethyl acetate (200 ml). The combined ethyl acetate layer was washed with saturated brine (500 ml) and dried over Na_2SO_4 . Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by column chromatography to give the title compound (10.46 g).

IR ν (nujol) cm^{-1} ; 3410, 1717, 1624, 1506.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.22 (3H, t, $J=7.0\text{Hz}$), 3.06 (2H, d, $J=5.0\text{Hz}$), 3.66 (1H, t, $J=5.0\text{Hz}$), 3.78 (2H, s), 3.90 (2H, s), 4.13 (2H, q, $J=7.0\text{Hz}$), 6.37 (1H, d, $J=2.0\text{Hz}$), 6.56 (1H, dd, $J=2.0, 8.4\text{Hz}$), 6.92 (1H, d, $J=8.4\text{Hz}$), 7.20-7.50 (5H, m).

Reference Example 5

2-(N-tert-Butoxycarbonyl-N-methylamino)ethyl methanesulfonate

[0167]

(1) 2-(Methylamino)ethanol (3.5 ml) was dissolved in tetrahydrofuran (150 ml). Thereto was added di-tert-butyl dicarbonate (12.5 g) and the mixture was stirred at room temperature for 20 min. Tetrahydrofuran was evaporated under reduced pressure. The obtained residue was purified by column chromatography to give 2-(N-tert-butoxycarbonyl-N-methylamino)ethanol (6.35 g).

2-(N-tert-Butoxycarbonyl-N-methylamino)ethanol:

IR ν (neat) cm^{-1} ; 3423, 2976, 2934, 2882, 1674.

$^1\text{H-NMR}$ (CDCl_3) δ (ppm);

1.43 (9H, s), 2.89 (3H, s), 3.34 (2H, t, $J=5.8\text{Hz}$), 3.67 (2H, t, $J=5.8\text{Hz}$), 4.00-6.00 (1H, br).

(2) The compound (505 mg) obtained in (1) above was dissolved in methylene chloride (20 ml). Thereto were added triethylamine (0.5 ml) and methanesulfonyl chloride (0.25 ml) and the mixture was stirred at room temperature for 1 hr. Methylene chloride (30 ml) was added to the mixture, and the mixture was washed with saturated brine (20 ml) and dried over Na_2SO_4 . Methylene chloride was evaporated under reduced pressure to give the title compound (720 mg).

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) ;

1.46 (9H, s), 2.94 (3H, s), 3.01 (3H, s), 3.54 (2H, t, J=5.5Hz), 4.33 (2H, t, J=5.5Hz).

Reference Example 6

1-(2-Bromoethyl)indoline

[0168] To 1,2-dibromoethane (58.0 ml) were added indoline (5.0 g) and triethylamine (28.7 ml), and the mixture was stirred at 90°C for 2 hr. Ethyl acetate (200 ml) was added to the reaction mixture, and the mixture was washed with saturated brine (400 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by column chromatography to give the title compound (4.09 g).

IR ν (neat) cm⁻¹ ; 2924, 2845, 1607, 1489.

¹H-NMR (CDCl₃) δ (ppm) ;

2.99 (1H, t, J=8.4Hz), 3.45 (1H, t, J=8.4Hz), 3.49 (4H, s), 6.40-6.75 (2H, m), 6.90-7.20 (2H, m).

Reference Example 7

1-(2-Bromoethyl)-6-methoxycarbonylindoline

[0169] To 1,2-dibromoethane (15.7 ml) were added 6-methoxycarbonylindoline (2.41 g) and triethylamine (7.8 ml), and the mixture was stirred at 90°C for 2 hr. Ethyl acetate (150 ml) was added to the reaction mixture, and the mixture was washed with saturated brine (300 ml) and dried over Na₂SO₄. Ethyl acetate was evaporated under reduced pressure. The obtained residue was purified by column chromatography to give the title compound (1.71 g).

IR ν (neat) cm⁻¹ ; 1713, 1611, 1499.

¹H-NMR (CDCl₃) δ (ppm) ;

3.03 (1H, t, J=8.4Hz), 3.53 (1H, t, J=8.4Hz), 3.53 (4H, s), 3.88 (3H, s), 6.00-6.20 (2H, m), 7.39 (1H, dd, J=1.5, 7.8Hz).

Experimental Example 1 Hypoglycemic action (method A)

[0170] Blood was drawn under a non-fasting state from the tail vein of male KK-A^y mice, which were spontaneously diabetic models, had developed diabetes due to insulin resistance and which showed hyperglycemia and hyperinsulinemia. The glucose level in the plasma was measured using a commercially available assay kit (glucose CII-test WAKO, Wako Pure Chemical Industries, Ltd.). The mice were grouped (5 mice per group) into a control group and an administration group, such that the mean and the standard deviation of the glucose level in the plasma of each group were nearly the same. Each test compound was suspended or dissolved in 5% gum arabic solution and administered orally to the administration group for 4 consecutive days from the next day. The 5% gum arabic solution was orally administered to the control group. Blood was drawn from the tail vein under a non-fasting state about 24 hours after the final administration and the glucose level in the plasma was measured. The decrease in the plasma glucose level was calculated from the following formula. The results are shown in Table 1.

Decrease (%) in plasma glucose level = [(mean plasma glucose level of control group - mean plasma glucose level of test compound administration group) / mean plasma glucose level of control group] × 100

Table 1 Hypoglycemic action (method A)

Test compound	Dose (mg/kg)	Decrease (%) in plasma glucose level	Test compound	Dose (mg/kg)	Decrease (%) in plasma glucose level
Example 2	10	38.3	Example 40	30	46.3
	30	60.6	Example 41	30	11.1
Example 3	30	11.8	Example 42	10	27.6
Example 5	10	34.1		30	49.6
	30	43.4	Example 43	30	27.5
Example 6	10	10.7	Example 44	30	48.0
	30	12.2	Example 45	30	12.6
Example 7	10	11.4	Example 46	30	28.6
	30	17.4	Example 48	30	16.1
Example 9	30	20.3	Example 49	30	11.1
Example 12	10	12.0	Example 50	30	11.4
Example 16	30	34.5	Example 51	30	20.0
Example 18	30	39.7	Example 52	30	26.3
Example 21	30	43.4	Example 53	30	13.6
Example 23	10	24.3	Example 54	30	10.8
	30	42.9	Example 55	30	13.9
Example 32	30	36.4	Example 58	30	19.1
Example 33	30	38.8	Example 67	30	10.6
Example 34	30	23.5	Example 69	30	12.5
Example 35	30	22.1	Example 70	30	15.2
Example 36	30	28.4	Example 71	30	10.5
Example 37	30	11.1	Example 72	30	15.9
Example 38	30	28.0	Example 73	30	14.8
Example 39	30	30.7			

Experimental Example 2 Hypoglycemic action (method B)

[0171] Blood was drawn under a non-fasting state from the tail vein of male KK-A^y mice, which were spontaneously diabetic models, had developed diabetes due to insulin resistance and which showed hyperglucemia and hyperinsulinemia. The glucose level in the plasma was measured using a commercially available assay kit (glucose CII-test WAKO, Wako Pure Chemical Industries, Ltd.). The mice were grouped (5 mice per group) into a control group and an administration group, such that the mean and the standard deviation of the glucose level in the plasma of each group were nearly the same. Each test compound was admixed in a proportion of 0.1 w/w% with a powder feed (CE-2, Clea) and the feed mixture was fed to the administration group for 4 days from the next day. A regular powder feed was fed to the control group. At day 5, blood was drawn from the tail vein under a non-fasting state and the glucose level in the plasma was measured. The decrease in the plasma glucose level was calculated from the following formula. The results are shown in Table 2.

Decrease (%) in plasma glucose level = [(mean plasma glucose level of control group - mean plasma glucose level of test compound administration group) / mean plasma glucose level of control group] × 100

Table 2 Hypoglycemic action (method B)	
Test compound	Decrease (%) in plasma glucose level
Example 2	69.2
Example 10	42.4

Experimental Example 3 hypotriglyceridemic action

[0172] Blood was drawn under a non-fasting state from the tail vein of male KK-A^y mice, which were spontaneously diabetic models, had developed diabetes due to insulin resistance, and which showed hyperglycemia and hyperinsulinemia. The triglyceride level in the plasma was measured using a commercially available assay kit (triglyceride G-test WAKO, Wako Pure Chemical Industries, Ltd.). The mice were grouped (5 mice per group) into a control group and an administration group, such that the mean and the standard deviation of the triglyceride level in the plasma of each group were nearly the same. Each test compound was suspended or dissolved in 5% gum arabic solution and administered orally to the administration group for 4 consecutive days from the next day. The 5% gum arabic solution was orally administered to the control group. Blood was drawn from the tail vein under a non-fasting state about 24 hours after the final administration and the triglyceride level in the plasma was measured. The decrease in the plasma triglyceride level was calculated from the following formula. The results are shown in Table 3.

Decrease (%) in plasma triglyceride level = [(mean plasma triglyceride level of control group - mean plasma triglyceride level of test compound administration group) / mean plasma triglyceride level of control group] × 100

Table 3

hypotriglyceridemic action					
Test compound	Dose (mg/kg)	Decrease (%) in plasma triglyceride level	Test compound	Dose (mg/kg)	Decrease (%) in plasma triglyceride level
Example 2	10	39.5	Example 38	30	30.0
	30	54.3	Example 39	30	15.6
Example 4	10	14.3	Example 40	30	36.1
Example 5	30	30.9	Example 42	30	31.6
Example 11	10	11.2	Example 43	30	20.9
Example 12	10	22.9	Example 44	30	19.1
Example 16	30	19.8	Example 49	30	17.0
Example 18	30	45.5	Example 50	30	35.7
Example 21	30	24.4	Example 51	30	10.7

Table 3 (continued)

hypotriglyceridemic action					
Test compound	Dose (mg/kg)	Decrease (%) in plasma triglyceride level	Test compound	Dose (mg/kg)	Decrease (%) in plasma triglyceride level
Example 23	10	30.4	Example 52	30	26.6
	30	50.4	Example 56	30	14.0
Example 32	30	32.5	Example 58	30	24.7
Example 33	30	42.0	Example 59	30	13.7
Example 34	30	17.3	Example 69	30	15.2
Example 35	30	12.4	Example 72	30	24.0
Example 36	30	13.7	Example 73	30	15.4

Experimental Example 4 Hypoglycemic action and hypoinsulinemic action in insulin resistant diabetic mouse

[0173] The insulin resistance improving action was examined in KK-A^y mice, which were spontaneously diabetic models, had developed diabetes due to insulin resistance, and which showed hyperglycemia and hyperinsulinemia. Blood was drawn from the tail vein of 12-week-old male KK-A^y mice under a non-fasting state, and the glucose level in the plasma was measured using a commercially available assay kit (glucose CII-test WAKO, Wako Pure Chemical Industries, Ltd.). The mice were grouped (5 mice per group) into a control group and an administration group, such that the mean glucose level in the plasma and the average body weight of each group, as well as the standard deviation thereof were nearly the same. The test compound (10 mg/kg) was suspended in 5% aqueous gum arabic solution and orally administered once a day to the administration group for 4 days from the next day. The 5% aqueous gum arabic solution was orally administered to the control group. Blood was drawn from the tail vein under a non-fasting state 24 hours after the final administration. The concentrations of plasma glucose and plasma insulin were measured. The results are shown in Table 4.

[0174] As a result, each test compound (10 mg/kg) lowered the glucose level in plasma and also decreased the insulin concentration in plasma. This means that the test compound lowered the plasma glucose level not by insulin secretion but by insulin sensitivity potentiating action (insulin resistance improving action), thereby to improve hyperinsulinemia.

Table 4

Hypoglycemic action and Hypoinsulinemic action			
Test compound	Dose of test compound (mg/kg)	Plasma glucose (mg/dl)	Insulin (ng/ml)
Control	0	507	43
Example 2	10	313	26
Example 23	10	382	28
Example 32	10	402	30
Example 42	10	308	27

Experimental Example 5 Triglyceride accumulation promoting action in 3T3-L1 cells

[0175] A culture medium of 80% confluent 3T3-L1 cells was removed and the cells were dissociated with 0.25% trypsin-EDTA solution. 5% FBS-DMEM (the same amount as the removed medium) was added and the obtained cell suspension was centrifuged at 25°C, 100×g for 1 min to give cell sediment and the supernatant was removed. The cells were re-suspended in a suitable amount of 5% FBS-DMEM medium and the cells were counted. 5% FBS-DMEM medium was added to adjust the concentration to 1×10⁵ cells/ml and the mixture was dispensed by 1 ml to a 24 well plate. The cells were cultured with aeration of 5% CO₂ at 37°C for 2 days. Upon confirmation of a post-confluent state, the culture supernatant was changed to a medium containing 0.5 mM-IBMX, and the cells were cultured for 2 days. Then, the medium was changed to a medium containing 10 ng/ml insulin and 10⁻⁷ M test compound and the cells were

cultured further for 4 days. After removing the culture supernatant, the cells were lysed in a 0.1% SDS solution and the triglyceride amount was measured. The triglyceride accumulation (%) due to the potentiation of insulin activity by the test compound was calculated from the following formula. The results are shown in Table 5.

$$\frac{[(\text{triglyceride amount with addition of test compound} - \text{triglyceride amount of control}) / \text{triglyceride amount of control}] \times 100}{}$$

Table 5 Triglyceride accumulation promoting action

Test compound	Triglyceride accumulation (%)	Test compound	Triglyceride accumulation (%)
Example 2	260.4	Example 33	222.5
Example 5	233.0	Example 37	277.3
Example 8	275.5	Example 39	258.0
Example 16	288.9	Example 40	231.0
Example 21	284.9	Example 42	193.6
Example 23	214.2	Example 62	327.8
Example 32	181.2		


Effect of the Invention

[0176] The heterocyclic compound [I] and a pharmaceutically acceptable salt of the present invention show a superior hypoglycemic action, a blood hypolipidemic action, an insulin resistance-improving action and a PPAR activating action and are useful as hypoglycemic agents; hypolipidemic agents, insulin resistance improvers, therapeutic agents of diabetes, therapeutic agents of diabetic complications, glucose tolerance improvers, anti-atherosclerosis agents, anti-obesity agents, antiinflammatory agents, agents for the prophylaxis or treatment of PPAR-mediated disease and agents for the prophylaxis or treatment of syndrome X. Therefore, the heterocyclic compound [I] and a pharmaceutically acceptable salt of the present invention are useful for the prophylaxis or treatment of diabetes, diabetic complications, hyperlipidemia, atherosclerosis, hyperglycemia, diseases caused by insulin resistant impaired glucose tolerance and diseases caused by insulin resistance, obesity, inflammation, PPAR-mediated disease and syndrome X. The heterocyclic compound [I] of the present invention has a structure completely different from the structure of the compounds heretofore used as an active ingredient of conventional insulin resistance improvers. By the provision of this compound, hypoglycemic agents, hypolipidemic agents, insulin resistance improvers, therapeutic agents of diabetes, therapeutic agents of diabetic complications, glucose tolerance improvers, anti-atherosclerosis agents, anti-obesity agents, anti-inflammatory agents, agents for the prophylaxis or treatment of PPAR-mediated disease and agents for the prophylaxis or treatment of syndrome X come to have a wide variety, from which a desired agent can be freely selected.

[0177] This application is based on patent application Nos. 345543/1999 and 295108/2000 filed in Japan, the contents of which are hereby incorporated by reference.

Claims

1. A heterocyclic compound of the formula [I]



wherein

- R¹ is hydrogen atom or lower alkyl,
 R² is hydrogen atom, alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent, alkenyl optionally having a substituent, alkynyl optionally having a substituent, heterocyclic alkyl optionally having a substituent or -COR⁴
 5 wherein R⁴ is hydrogen atom, alkyl optionally having a substituent, aryl optionally having a substituent, alkenyl optionally having a substituent, arylalkyl optionally having a substituent or heterocyclic residue optionally having a substituent
 R³ is hydrogen atom, lower alkyl or lower alkoxy,
 A is a single bond or >N-R⁵
 10 wherein R⁵ is hydrogen atom or loweralkyl,
 B is lower alkylene, and
 Y is aryl optionally having a substituent or an aromatic heterocyclic residue optionally having a substituent

or a pharmaceutically acceptable salt thereof.

- 15 2. The heterocyclic compound of claim 1, wherein, in the formula [I],

R¹ is hydrogen atom or lower alkyl,
 R² is hydrogen atom, alkyl optionally having a substituent, cycloalkyl, cycloalkylalkyl, aryl optionally having a substituent, arylalkyl optionally having a substituent or -COR⁴ wherein R⁴ is hydrogen atom, alkyl optionally having a substituent, aryl optionally having a substituent or arylalkyl optionally having a substituent,
 20 R³ is hydrogen atom, lower alkyl or lower alkoxy,
 A is a single bond or >N-R⁵ wherein R⁵ is hydrogen atom or lower alkyl,
 B is lower alkylene, and
 25 Y is aryl optionally having a substituent or an aromatic heterocyclic residue optionally having a substituent,

or a pharmaceutically acceptable salt thereof.

- 30 3. The heterocyclic compound of claim 1, wherein, in the formula [I],

R¹ is hydrogen atom or lower alkyl,
 R² is hydrogen atom, alkyl, cycloalkylalkyl, arylalkyl optionally having a substituent, alkenyl, alkynyl, heterocyclic alkyl or
 -COR⁴ wherein R⁴ is alkyl, alkenyl or aryl,
 35 R³ is hydrogen atom or lower alkoxy,
 A is a single bond or >N-R⁵ wherein R⁵ is lower alkyl,
 B is lower alkylene, and
 Y is aryl or an aromatic heterocyclic residue optionally having a substituent,

40 or a pharmaceutically acceptable salt thereof.




4. The heterocyclic compound of claim 1, wherein, in the formula [I],

R¹ is hydrogen atom or lower alkyl,
 45 R² is hydrogen atom, alkyl, cycloalkylalkyl, arylalkyl optionally having a substituent or -COR⁴ wherein R⁴ is alkyl or aryl,
 R³ is hydrogen atom,
 A is a single bond or >N-R⁵ wherein R⁵ is lower alkyl,
 B is lower alkylene, and
 50 Y is an aromatic heterocyclic residue optionally having a substituent,

or a pharmaceutically acceptable salt thereof.

- 55 5. The heterocyclic compound of claim 1, wherein, in the formula [I],

Y-A- is



30 wherein

R^A is isopropyl or tert-butyl,



R^B is isopropyl or tert-butyl,


R^C is isopropyl, tert-butyl, phenyl, thiophen-2-yl, 2-methylpropenyl, 3-butenyl, cyclopropyl, 1-butenyl or 2,2-dimethylpropyl,

35 or a pharmaceutically acceptable salt thereof.


6. The heterocyclic compound of claim 1, wherein, in the formula [I],

40 Y-A- is





or



wherein





R^A is isopropyl or tert-butyl,

R^B is isopropyl or tert-butyl,


R^C is isopropyl, tert-butyl, phenyl, thiophen-2-yl, 2-methylpropenyl or 3-butenyl,

or a pharmaceutically acceptable salt thereof.

7. The heterocyclic compound of claim 1, wherein, in the formula (I),
Y-A- is




or



or a pharmaceutically acceptable salt thereof.

8. The heterocyclic compound of claim 1, wherein, in the formula [I],
Y-A- is



or a pharmaceutically acceptable salt thereof.

9. The heterocyclic compound of claim 1, wherein the heterocyclic compound of the formula [I] is any of the following compounds (1) to (67):

- (1) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (2) 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (3) 2-acetyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (4) 2-methyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (5) 2-hexanoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (6) 2-hexyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (7) 2-isobutyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (8) 2-cyclohexylmethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (9) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(3-phenylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (10) 2-benzoyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (11) 2-benzyl-7-[2-(N-methyl-N-(pyridin-2-yl)amino)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (12) 2-benzyl-7-[2-(5-ethyl-pyridin-2-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (13) 2-benzyl-7-[2-(indolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (14) ethyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (15) methyl 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (16) 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (17) ethyl 2-(4-methoxybenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (18) 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (19) ethyl 2-(4-methylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
- (20) 2-benzyl-7-[2-(6-carboxyindolin-1-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (21) 2-(4-fluorobenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (22) 2-(2,2-dimethylpropionyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (23) 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (24) 2-benzyl-7-[2-(5-methyl-2-tert-butyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (25) 2-benzyl-7-[2-(5-methyl-2-(thiophen-2-yl)oxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (26) 2-benzyl-7-[2-(5-methyl-2-isopropylloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (27) 2-butyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (28) 2-benzyl-7-[2-[5-methyl-2-(2-methylpropenyl)oxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (29) 2-benzyl-7-[2-(3-butenyl)-5-methyloxazol-4-yl]ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (30) 2-allyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (31) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(2-propynyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (32) 2-(2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (33) 2-benzyl-7-[(indolin-3-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
- (34) 2-(3-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,

- (35) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-pentanoyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (36) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(4-pentenoyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 5 (37) 2-(3-methyl-2-butenoyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (38) 2-(3,3-dimethylbutyryl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 10 (39) 2-benzyl-7-methoxy-6-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3RS)-carboxylic acid,
 (40) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(pyridin-2-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (41) 2-benzyl-7-(3-methyl-3-phenylbutoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (42) 2-benzyl-7-(3,3-dimethyl-4-phenylbutoxy)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 15 (43) 2-benzyl-7-(2-isopropylbenzoxazol-6-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (44) 2-benzyl-7-(2-tert-butylbenzoxazol-6-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (45) 2-benzyl-7-(2-tert-butylbenzoxazol-5-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (46) 7-(2-tert-butylbenzoxazol-6-yl)methoxy-2-(2,2-dimethylpropyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 20 (47) 2-benzyl-7-(2-isopropylbenzoxazol-5-yl)methoxy-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (48) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(pyridin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (49) 7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-[(pyridin-2-yl)carbonyl]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 25 (50) methyl 2-benzyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
 (51) 2-benzyl-7-[2-(2-cyclopropyl-5-methyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (52) 2-(3-methyl-2-butenyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 30 (53) 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-tert-butyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (54) 2-benzyl-7-[2-(1-butenyl)-5-methyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 35 (55) 2-benzyl-7-[2-(2,2-dimethylpropyl)-5-methyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (56) ethyl 2-(2,2-dimethylpropyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylate,
 (57) 7-(benzofuran-2-ylmethoxy)-2-benzyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 40 (58) 2-isobutyryl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (59) 7-[2-(benzofuran-2-yl)ethoxy]-2-benzyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (60) 7-[2-(5-ethylpyridin-2-yl)ethoxy]-2-hexanoyl-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (61) 2-carboxymethyl-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 45 (62) 2-[3-(methoxycarbonyl)propionyl]-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (63) 2-[3-(ethoxycarbonyl)propyl]-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 50 (64) 2-benzyl-6-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3RS)-carboxylic acid,
 (65) 2-(3-acetylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid,
 (66) 2-(2-acetylbenzyl)-7-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid, and
 55 (67) 2-benzyl-7-[(5-methyl-2-phenyloxazol-4-yl)methoxy]-1,2,3,4-tetrahydroisoquinoline-(3S)-carboxylic acid, or a pharmaceutically acceptable salt thereof.

10. The heterocyclic compound of claim 9, wherein the heterocyclic compound of the formula [I] is any of the above-mentioned compounds (1) to (47), or a pharmaceutically acceptable salt thereof.
11. The heterocyclic compound of claim 9, wherein the heterocyclic compound of the formula [I] is any of the above-mentioned compounds (1) to (21), or a pharmaceutically acceptable salt thereof.
12. A pharmaceutical composition comprising the heterocyclic compound of any of claim 1 to claim 11 or a pharmaceutically acceptable salt thereof.
13. A pharmaceutical agent comprising the heterocyclic compound of any of claim 1 to claim 11 or a pharmaceutically acceptable salt thereof, which is selected from the group consisting of a hypoglycemic agent, a hypolipidemic agent, an insulin resistance improver, a therapeutic agent of diabetes, a therapeutic agent of diabetic complication, a glucose tolerance improver, an anti-atherosclerosis agent, an anti-obesity agent, an antiinflammatory agent, an agent for the prophylaxis or treatment of PPAR-mediated disease and an agent for the prophylaxis or treatment of syndrome X.
14. A hypoglycemic agent comprising the heterocyclic compound of any of claim 1 to claim 11 or a pharmaceutically acceptable salt thereof.
15. A hypolipidemic agent comprising the heterocyclic compound of any of claim 1 to claim 11 or a pharmaceutically acceptable salt thereof.
16. An insulin resistance improver comprising the heterocyclic compound of any of claim 1 to claim 11 or a pharmaceutically acceptable salt thereof.
17. A therapeutic agent of diabetic complication comprising the heterocyclic compound of any of claim 1 to claim 11 or a pharmaceutically acceptable salt thereof.
18. A therapeutic agent of diabetes comprising the heterocyclic compound of any of claim 1 to claim 11 or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP00/08464

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl. ⁷ C07D217/26, 401/12, 405/12, 413/12, 413/14, A61K31/472, 31/4725, A61P3/04, 3/06, 3/10, 9/10, 29/00 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl. ⁷ C07D217/26, 401/12, 405/12, 413/12, 413/14, A61K31/472, 31/4725, A61P3/04, 3/06, 3/10, 9/10, 29/00 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Jitsuyo Shinan Koho 1926-1992 Toroku Jitsuyo Shinan Koho 1994-1996 Kokai Jitsuyo Shinan Koho 1971-1992 Jitsuyo Shinan Toroku Koho 1996-2000 Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CA (STN), REGISTRY (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, 98/00403, A1 (ELI LILLY AND COMPANY), 08 January, 1998 (08.01.98), Full text & JP, 2000-515501, A	1-18
X	WO, 93/23378, A1 (WARNER-LAMBERT COMPANY), 25 November, 1993 (25.11.93), Full text & JP, 7-508266, A	1-6, 12, 13
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 24 January, 2001 (24.01.01)		Date of mailing of the international search report 06 February, 2001 (06.02.01)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)